


THE DUODECIMAL BULLETIN

Vol. 54_z (64_d), No. 1 • December 1206_z (2022_d)




Dozenal Card Games

Whole Number

74_z
(124_d)

ISSN 0046-0826_d



THE DOZENAL SOCIETY OF AMERICA

is a voluntary, nonprofit corporation, organized to conduct research and educate the public in the use of base twelve in calculations, mathematics, weights and measures, and other branches of pure and applied science.

As a nonprofit we depend upon the generosity of our members to continue operating. Basic membership is US\$12.00_d (US\$10.00_z). An annual subscription of US\$36.00_d (US\$30.00_z) entitles the member to delivery of paper copies of *The Duodecimal Bulletin*.

The DUODECIMAL BULLETIN
is an official publication of:

THE DOZENAL SOCIETY
OF AMERICA, INC.
3100 Central Ave, Suite 412
St Petersburg, FL 33712

FOUNDED: 1160_z (1944_d)

Official Website:
dozenal.org

Email us at:
contact@dozenal.org

Follow us on Twitter:
[@dozenal](https://twitter.com/dozenal)

Editor: John Volan
editor@dozenal.org

The Dozenal Society
of Great Britain:
dozenalsociety.org.uk

OFFICERS

| | |
|----------------|--------------------|
| BOARD CHAIR | Jay Schiffman |
| PRESIDENT | Graham Steele |
| VICE PRESIDENT | John Volan |
| SECRETARY | Michael De Vlieger |
| TREASURER | Jay Schiffman |

BOARD OF DIRECTORS

| | |
|--------------------|-------------------|
| Dean Canfield | SOUTHERN VA |
| Michael De Vlieger | ST. LOUIS, MO |
| John Impagliazzo | LONG ISLAND, NY |
| Paul Rapoport | HAMILTON, ON |
| Jay Schiffman | PHILADELPHIA, PA |
| Graham Steele | ST PETERSBURG, FL |
| Timothy Travis | FAYETTEVILLE, NC |
| John Volan | ASHLAND, MA |

DIRECTORS EMERITI

| | |
|----------------|-----------------|
| Donald Goodman | WOODBRIIDGE, VA |
| Jen Seron | NEW YORK, NY |
| Dan Simon | NEW YORK, NY |

DozensOnline Forum:
tapatalk.com/groups/dozensonline

The DSA does not endorse particular symbols for digits in alternate bases. Authors are free to propose their own symbols. However, unless otherwise specified, this publication defaults to the Pitman numerals for the digits ten (ε) and eleven (E).

The DSA does not endorse any particular scheme for disambiguating the base of a number. Authors are free to propose their own notations. However, as a default this publication will annotate the base of any number containing more than one digit using a single-letter subscript: d for Decimal, z for doZenal, x for heXadecimal, o for Octal, b for Binary, etc. By default, numbers are annotated individually, although whole numeric expressions in parentheses may be given a single annotation applying to all of their components. A tabular display may omit annotations, as long as the base used by the entire display is indicated.

The Duodecimal Bulletin, Volume Five Dozen Four, Number One, Whole Number Ten Dozen Four. Copyright © 1206_z (2022_d), The Dozenal Society of America.

The **DUODECIMAL** **BULLETIN**

~ TABLE OF CONTENTS ~

EDITORIAL: IS A TURNAROUND IN THE CARDS? 4_z

NEW MEMBERS 6_z

DOZENAL CARD GAMES
by Paul Rapoport 8_z

TIME (AND TIME AGAIN)
by Paul Rapoport 10_z

THE ONLINE DOZENAL/DECIMAL SCIENTIFIC CALCULATOR
by Paul Rapoport 13_z

PRIMEL METROLOGY, PART II
by John Volan 16_z

SYSTEMATIC DOZENAL NOMENCLATURE SUMMARY 25_z

AN UNPUBLISHED MANUSCRIPT OF LEIBNIZ'S ON DUODECIMAL
by Lloyd Strickland 26_z

DOZENAL MAGIC SQUARES
by Jay L. Schiffman 31_z

POUNDS, SHILLINGS, PENCE: THE COMMONWEALTH'S DOZENAL CURRENCY
by Andon Epp 37_z

POETRY CORNER 42_z



See "The Opposed Principles" https://dozenal.org/drupal/sites_bck/default/files/db4a205_0.pdf
Image credit: Victoriano Izquierdo (Unsplash license)



Is a Turnaround *in the Cards*?

SINCE OUR LAST ISSUE, much in the world has changed. The COVID-03 epidemic¹ has been raging now for three years, forcing people of all walks of life to accept many adjustments we would never have imagined before. Perhaps the disease is attenuating, but it's not clear yet when, if ever, it will be past.

Meanwhile, the conflict devastating the Ukraine has been raging for the better part of a year, with as much of an uncertain outcome as the pandemic. An invasion of one country by another on the European continent, the first since World War II, was heretofore inconceivable in this age of nuclear stalemate. And yet, here it is.

Yet not all news is dire: As I write these words, the unmanned Artemis I spacecraft has just returned from its 21_z day mission to a picture-perfect splashdown, boosting hopes of human beings returning to the Moon for the first time in over four unquennia—and perhaps even going on to Mars within another unquennium.

Much has changed for the dozenal community as well. Donald Goodman III, after many fruitful years at the helm of the Dozenal Society of America, has elected to resign his position as President. We all thank him for his dedicated service and wish him the very best, for himself and his family. As an emeritus member of the Board of Directors, he remains a welcome friend to all in the DSA. Graham Steele, formerly our Vice President, has stepped in to fulfill Don's duties in the interim.

Meanwhile, the difficulties of the pandemic have prevented the DSA from holding an annual in-person meeting. On the other hand, like many during these times, we have learned to adapt by exploiting technology: The DSA Board has managed to meet several times virtually, via Zoom. We are looking into the possibility of using the same means to make the 1207_z Annual Meeting a virtual or hybrid event for the whole membership. We're also considering experimenting with other virtual events, such as presentations or discussion forums on focused topics of dozenal interest, that would be open to all via Zoom, and that could be recorded and made available online on our [dozenal.org](https://www.dozenal.org) website. To that end, all members who have not already done so are encouraged to join the DozensOnline forum,² as that is the best place to receive timely updates about DSA business.

Nevertheless, article submissions to the *Bulletin* remain robust. Paul Rapoport has graced us with no less than three articles detailing his indefatigable efforts to promote fun or useful online applications of dozenal. In fact, the cover article of the issue is his description of online solitaire card games he has developed which utilize dozenal numbers. He also gives us an update on his work to implement a dozenal wristwatch, as well as his project to develop an online dozenal/decimal scientific calculator. The latter has involved contract work from professional programmers, which Paul hired

¹So called, of course, because it began in the year 1203_z.

²<https://www.tapataalk.com/groups/dozensonline>

at his own considerable expense. Although he volunteered this as a contribution to the society, the *Bulletin* would like to appeal to the membership to try out Paul's calculator, and if they find it useful, make whatever donation to the DSA they can and earmark it to defray Paul's cost.

My own contribution to this issue is another installment about the Primel metrology I introduced in the last issue, this time covering the topic of angular mechanics. Perhaps considering how to adapt a coherent dozenal-metric metrology to the physics of objects turning can help us all contemplate how to turn things around in these times.

Professor Lloyd Strickland of Manchester Metropolitan University in Great Britain has discovered a remarkable bit of manuscript by none other than the mathematician and philosopher Gottfried Wilhelm Leibniz. It shows a fascinating glimpse at how Leibniz, more than 230_z years ago, was analyzing the properties of the duodecimal base—and doing so, of course, in Latin, per the custom of scientists of that era. This historically interesting manuscript has never been published before, but Professor Strickland thought of us, and graciously offered to give the *Bulletin* first dibs at bringing it to light.


Our own Professor Jay Schiffman, DSA Treasurer, offers another of his mathematical morsels, using both decimal and dozenal base to analyze third-order magic squares, including magic squares containing primes.

One of our younger members, Andon Epp, who came to an Annual Meeting with his parents a few years ago, has written a nice retrospective on the old British pound-shillings-pence currency, complementing it with research on what dozenalists have more recently speculated concerning what a truly dozenal currency might look like. We're encouraged for the future that such thoughtful youths are stepping up to participate. Our recently departed emeritus member Gene Zirkel, longest-serving by far, got his own start in the society writing an article for the *Bulletin* nearly six dozen years ago, when he was not much older than Andon is now.

In fact, Michael de Vlieger, DSA Secretary and former editor of the *Bulletin*, is right now in the process of putting together another issue, to be published shortly after this one, which will be completely devoted as a tribute to Gene Zirkel. Look for that to arrive early in 1207_z.

So whatever is in the cards, let this be my holiday card to you all: Here's wishing that this season finds all of our membership well and happy, and enjoying among other things the elegance of dozenal arithmetic, during this dozenth month of 1206_z. However life is treating you, may things turn around to a brighter future in 1207_z! 🍀

NEW MEMBERS




SINCE the last issue, we've seen continuing growth in our membership. Our rolls have gone up by nearly another third.

Joining is easy! You can do it electronically, at the DSA's website: dozenal.org. Just click the **Join Us!** button on the top right. A basic membership is free. For a donation of \$30_z (\$36_d) per year, members can subscribe to receive hard copies of the *Bulletin* as they are published. (Subscribing members are highlighted in red below. The electronic version is free to all members.) The DSA Board would like to invite all of our members, new and old, to come to our annual meetings. We'd love to meet you all! Feel free to email editor@dozenal.org your ideas for future *Bulletin* articles. Make sure to visit the DozensOnline Forum at <https://www.tapataalk.com/groups/dozensonline/> to chat with us about all subjects related to dozenal and to watch out for announcements to all members.

| | |
|---|---|
| 76 _z (1134 _d) Wolfram Harper | 823 _z (1179 _d) Christian G. Wampera |
| 767 _z (1135 _d) Kelly Jonathan Christoffersen | 824 _z (1180 _d) Karan Pundir |
| 768 _z (1136 _d) Kylie Gorney | 825 _z (1181 _d) Joseph Nudell |
| 768 _z (1139 _d) Declan Powell | 826 _z (1182 _d) Connor Robert Cochran |
| 781 _z (1141 _d) Jerzy K. Malewicz | 827 _z (1183 _d) Douglas James Butner |
| 782 _z (1142 _d) Justin Lee McCall | 828 _z (1184 _d) Noah Michael Howard |
| 783 _z (1143 _d) Gage Trevor Norton | 829 _z (1185 _d) Ethan John Koetsier |
| 784 _z (1144 _d) Soumyabrata Bhattacharya | 829 _z (1186 _d) Neil A |
| 785 _z (1145 _d) Alexander Channing Rowland | 829 _z (1187 _d) Jon Bennett |
| 786 _z (1146 _d) Matt Pope | 830 _z (1188 _d) Phunsook Wangdoo |
| 787 _z (1147 _d) John F. Schilke, MD | 831 _z (1189 _d) James Jacob Wilson |
| 788 _z (1148 _d) Levi C Perrine | 832 _z (1190 _d) Meredith K. Saylor |
| 789 _z (1149 _d) Kenneth P Wolcott | 833 _z (1191 _d) Steele Knudson |
| 789 _z (1150 _d) Grant David Meadors | 834 _z (1192 _d) Mario Butera |
| 789 _z (1151 _d) Christian Czarnowsky | 835 _z (1193 _d) Ricky Guerin |
| 800 _z (1152 _d) Luke Josiah Feezell | 836 _z (1194 _d) John Bender |
| 801 _z (1153 _d) John Walter | 837 _z (1195 _d) Ty Jacob Gozzard |
| 802 _z (1154 _d) André M. Tristant | 838 _z (1196 _d) Johan Kovacs |
| 803 _z (1155 _d) Brian Lee | 839 _z (1197 _d) Chris Gale |
| 804 _z (1156 _d) Davide Dalco | 839 _z (1198 _d) Peter John Wright |
| 805 _z (1157 _d) Frank Hunleth | 840 _z (1200 _d) Johnathon Warren Howell |
| 806 _z (1158 _d) Megan Hunleth | 842 _z (1202 _d) Paula G Evans |
| 807 _z (1159 _d) Parker Declan Schultz | 843 _z (1203 _d) Saksham Chakrawal |
| 808 _z (1160 _d) Robin Vahimian | 844 _z (1204 _d) Santosh Paripelli |
| 809 _z (1161 _d) Erick K Hoegberg | 845 _z (1205 _d) Jerry Barnish |
| 809 _z (1162 _d) Ramsay Midwood | 846 _z (1206 _d) Jason Yust |
| 809 _z (1163 _d) Shehrose Mian | 847 _z (1207 _d) Terence R Wulff |
| 810 _z (1164 _d) Mark Hollifield | 848 _z (1208 _d) Dylan James Haugh-Ewald |
| 811 _z (1165 _d) Thomas Rockwell Grose | 849 _z (1209 _d) Todd Call Partridge |
| 812 _z (1166 _d) Nathan Richard Krause | 849 _z (1210 _d) Sebastian "Pszczola" Dryl |
| 813 _z (1167 _d) Christopher Michael Hackwith | 850 _z (1212 _d) Joseph Dean |
| 814 _z (1168 _d) Harold Brosowsky | 851 _z (1213 _d) Courtney Brianne Harper |
| 815 _z (1169 _d) Rafal Studziński | 852 _z (1214 _d) Fara Tyler |
| 816 _z (1170 _d) Eric Lachapelle | 853 _z (1215 _d) Austin Tyler Sawyer |
| 817 _z (1171 _d) Nando Sanchez | 854 _z (1216 _d) Richard Frank Knackstedt Dillon |
| 818 _z (1172 _d) Jane Kim | 855 _z (1217 _d) Dylan Lupo |
| 819 _z (1173 _d) Andrew Moorman | 856 _z (1218 _d) Matthew Joseph Loewen |
| 819 _z (1174 _d) Maxime Gosso | 857 _z (1219 _d) Pi Francis |
| 819 _z (1175 _d) James Maxwell Bal | 858 _z (1220 _d) Tom Grogan |
| 820 _z (1176 _d) Matias Silva Mella | 859 _z (1221 _d) Bastien Magnetoscope |
| 821 _z (1177 _d) Evan Scott | 859 _z (1222 _d) Alexis Mekenzie Caplener |
| 822 _z (1178 _d) Moses Robert McKinney | 859 _z (1223 _d) Andrei Khomka |

860_z (1224_d) Patrick Adelman
 861_z (1225_d) Francis G
 862_z (1226_d) Mark von Germeten
 863_z (1227_d) Paul Henry Tegelaar
 865_z (1229_d) Raymond N Phillips
 866_z (1230_d) KellyJo Tarbell
 867_z (1231_d) Loretta D Sartin
 869_z (1233_d) John Rakauskas
 867_z (1234_d) John M Landry
 868_z (1235_d) Gregory Flagg
 870_z (1236_d) Scoot Black
 871_z (1237_d) Alexander Joseph Hostetler
 872_z (1238_d) William D Elkins
 873_z (1239_d) Edmund John Meurer
 874_z (1240_d) David McCarroll
 875_z (1241_d) Melvin Randall Moon
 876_z (1242_d) Cheese J Hasselberger
 877_z (1243_d) David McCarroll
 879_z (1245_d) Salvatore P Spatafore
 877_z (1246_d) Terese Brasile
 878_z (1247_d) Mark Jacob Toney
 880_z (1248_d) Willow Anne Starc
 881_z (1249_d) John Richard Kennedy
 882_z (1250_d) Peter K Rotich
 883_z (1251_d) Asuquo Levy Eyo
 885_z (1253_d) Arielle Marq
 886_z (1254_d) Matthew Hiscoe
 887_z (1255_d) Aaron Hopus
 888_z (1256_d) Lauren E. Haupt
 889_z (1257_d) Don T Ford
 887_z (1258_d) Tess Andres
 888_z (1259_d) Christopher John Alford
 890_z (1260_d) Richard D Hagedorn
 891_z (1261_d) C Spencer Chamberlain
 892_z (1262_d) William R Bruce
 893_z (1263_d) Keith T Vezeau
 894_z (1264_d) Kyle Licht
 895_z (1265_d) Mahyar Sahraei
 896_z (1266_d) Jayson Quilantan
 897_z (1267_d) Stephen Robinson
 898_z (1268_d) Robert Richard Muller
 899_z (1269_d) Brian Lee
 870_z (1272_d) John Michael Sefel
 871_z (1273_d) Vafaiei Radin
 872_z (1274_d) Amanda Sue Yale
 873_z (1275_d) Joshua David Cain
 874_z (1276_d) Sean P Tarrant
 875_z (1277_d) Gabriel Michael Gervais
 876_z (1278_d) Lanice Angelina Soulier
 877_z (1279_d) Todd Richardson
 878_z (1280_d) Brian Reynolds
 879_z (1281_d) Daniela Michelle Large
 877_z (1282_d) **Mikhael Etan Lyvers**
 878_z (1283_d) Johan Pattyn
 882_z (1286_d) Damien Boizard
 883_z (1287_d) Brian Schwartz
 884_z (1288_d) **Richard Dreselly**
 885_z (1289_d) Dustin Swan
 886_z (1290_d) Ruben Jimenez Mejias
 887_z (1291_d) Mark Morra
 888_z (1292_d) Matthew Christopher Bartsh
 889_z (1293_d) Walter C Ruppman
 887_z (1294_d) Karen A Gogol
 888_z (1295_d) Marley Abigail Braly
 900_z (1296_d) Jonathan Paul Sharman
 901_z (1297_d) Justin Pecot
 902_z (1298_d) William Henderson
 903_z (1299_d) Matthew Wilkins
 904_z (1300_d) Hector Barraza-Atayde
 905_z (1301_d) Terry Leon Renfrow
 906_z (1302_d) Zach Williams
 907_z (1303_d) Mike Ropke
 908_z (1304_d) **John de Paul Hansen**
 909_z (1305_d) Merlin Lee
 907_z (1306_d) Sioux Renaissance Hart
 908_z (1307_d) Helen Reilly Pruitt
 910_z (1308_d) John Barry Bulger
 911_z (1309_d) Zachary L Standig
 913_z (1311_d) Chris Henderson
 914_z (1312_d) Robert L Stayton
 916_z (1314_d) Antoine Carolus
 917_z (1315_d) Zain Yousuf
 918_z (1316_d) Dylan Tyler Sanborn
 919_z (1317_d) Divyosmi Hindol Goswami
 917_z (1318_d) Jennifer A Smith
 918_z (1319_d) Kyle R Finley
 920_z (1320_d) Sean Chappell
 921_z (1321_d) Saffron Moosefull
 922_z (1322_d) Anandita Mishra
 923_z (1323_d) Isaiah Eathan Boggs
 924_z (1324_d) **Mark Atwood**
 925_z (1325_d) Joe Schneebaum
 926_z (1326_d) Brian Jacob Xiao
 927_z (1327_d) Benjamin Orman
 928_z (1328_d) Mikko Viikari
 929_z (1329_d) **John Morris**
 927_z (1330_d) Dennis P. Matthews
 928_z (1331_d) Nick Davis
 930_z (1332_d) Hyung Tae Carapeto Figur
 931_z (1333_d) **Paul Branden**
 932_z (1334_d) Tyler Greenwood
 934_z (1336_d) Max van Zwienen
 935_z (1337_d) Daniel Anthony Barry
 936_z (1338_d) Jeffery Randolph Murray
 937_z (1339_d) Robert Bonomo
 938_z (1340_d) Hunter Prueger
 939_z (1341_d) Giovanni A Orlando
 937_z (1342_d) Tanya Lee McGowan
 938_z (1343_d) Liam Roger Krueger
 940_z (1344_d) Jim Thiele
 941_z (1345_d) Hope Rayanne Halyalkar
 942_z (1346_d) Nick Hudak
 f 943_z (1347_d) Shawn O'Neil
 944_z (1348_d) Tim Plummer
 945_z (1349_d) Ronald W Hotchkiss
 946_z (1350_d) Megan O'Brien Walsh
 947_z (1351_d) Jenna Lee
 948_z (1352_d) Olga Alexandrovna Pyetukhova
 949_z (1353_d) Andrew Whelan Kirk
 947_z (1354_d) Adam Ammar Jabri
 947_z (1355_d) Randy Lane Payne
 950_z (1356_d) Alan Lane Causey
 951_z (1357_d) Roman Anatolievich Shaternikov
 952_z (1358_d) Aleksandr Alekseevich Adamov
 953_z (1359_d) Luis Pedro Insua
 954_z (1360_d) José Jaime Longoria
 955_z (1361_d) Martin Trillsam
 956_z (1362_d) Lea Lemieux
 957_z (1363_d) Kayley Doris Galbraith
 958_z (1364_d) William H Suphan
 959_z (1365_d) David M Troutman
 957_z (1366_d) Mo Mi
 958_z (1367_d) Kin Ayattem
 960_z (1367_d) Bila Aberra
 961_z (1368_d) Daniel Stock
 ■■■



DOZENAL CARD GAMES

by Paul Rapoport

WHAT CAN DOZENALISTS USE to attract other people to the dozenal base, without being stymied by the near universal use of decimal? It turns out one endeavor can expose people to dozenal numbers with relative ease while actually entertaining them: dozenal card games. Creating a dozenal deck is easy. In many games, a few adjustments in play or scoring make it possible for players to ignore decimal completely.

The traditional card deck comprises 52_d cards: four suits, each with ten numbered “pip cards” and three royal “face cards.” The simplest way to dozenalize a deck is to change the label on the ten-pip card from 10_d to 7 , then add an eleven-pip card labeled ε , and a dozen-pip card labeled 10_z . Voilà, a fully-dozenal deck of 50_z (60_d) cards.

Several years ago, the French chess historian Jean-Louis Cazaux created chess games on $12_d \times 12_d$ boards. He furthered his interest in twelve by creating a deck of cards that had pips up to 12_d . He named his deck *K6T*. It’s available at <https://www.thegamecrafter.com/games/k6t>.

Cazaux readily agreed to create a fully dozenal deck by changing the cards 10_d , 11_d , and 12_d to 7 , ε , and 10_z . He also expanded his decimal deck in stages to make it even more dozenal: He added two new suits (moons and stars) to make a total of six, and added three new face cards, inspired by chess pieces, to make a total of six in each suit: jack (pawn), cavalier (knight), bishop, tower (rook), queen, king. Then he filled in more pip cards, now up to 24_d . The dozenal deck has followed, going up to 20_z pips. In the six suits, both decks include a 1 card in addition to the ace, plus a 0 card (intended mostly as a joker/wild card), and an infinity card (whose use has yet to be determined). Two additional blank cards round out the total in either deck to 148_z (200_d) cards.

Most card players will wonder why so many cards, when the traditional deck has sufficed for centuries. I suggest that more cards allow the invention of new games that are at least as engaging and fun as traditional games, while offering whole new perspectives to the players. At any rate, players may readily choose how many cards to use for any given game.


In 1203–1204_z (2019–2020_d), I produced three dozenal solitaire card games for physical play, as well as for online play at <https://games.dozenal.ca/solitaire/pyramid/>. Their dozenal aspect is vital. The purpose of these games is to teach or reinforce dozenal addition and multiplication in appealing ways.

PYRAMID ADDITION


The first game, Pyramid Addition, is similar to the common pyramid solitaire but with a few twists. The dozenal deck size may vary from 40_z to 110_z . In the smallest,

the pips run from A to 9; in the largest, from 0 to 16_z, with both the 1 card and the ace, the ace being the highest card.

The object of the game is simply to empty the tableau by removing pairs of exposed cards that add up to a target. The player chooses the tableau size, a smaller one being much easier to win with than a larger. The example shown here uses the standard dozenal deck (50_z cards) with a small tableau (24_z), and target 13_z. To equal the target, the jack pairs with the 2, the queen with the ace. The king requires no pair, because it is equal to the target and because this configuration does not use the 0 card.



In the above game, the first moves will be to remove the 7 and 8, and the 5 with one of the 7's. The 7 of spades is the better choice, because then the 2 and the jack will be exposed for removal, followed by the remaining 7 and 5 as shown here.



When no pairs are removable, stock cards are turned over to enable forming a pair with an exposed card in the tableau. Points are totaled at the end of the game, awarded in relation to the size of the deck and tableau, and deducted in relation to how many actions the player has taken to win.

PYRAMID FACTORS

The second game is Pyramid Factors, whose deck is 100_z cards. Although the object of removing pairs is the same, this time the pairs must come from the same group. A “group” consists of numbers related in elementary theory as indicated below, but not already included in a group higher in the list:

- Powers of 2: 2, 4, 8, 14_z
- Multiples of 6: 6, 10_z, 16_z, K(20_z)
- Multiples of the first totative¹: 5, 7, 13_z
- Multiples of the second totative: 7, 12_z, B(19_z)
- Multiples of 3 and 4: 3, 9, C(18_z)


¹A *totative* has only 1 as a common factor with the base.

- Omega², alpha³, and their multiples: ε , 11_z , $T(17_z)$
- Primes: 15_z , $J(17_z)$, $Q(1\varepsilon_z)$
- One: each 1 is its own group, not paired with another card.

In addition, a pair must not come from adjacent suits.

An exception to coming from the same group is bonus pairs from the first two groups that form a product of 10_z , 20_z , 30_z , 40_z , 60_z , 100_z . Most of these, e.g. 2×6 and 8×6 , have factors from both groups.

Below is a starting tableau of Pyramid Factors with 80_z cards, the maximum. In this game, rather than by portraits, the six face cards are symbolized by their related chess pieces.



Exposed for removal are a C and a 3, a 12_z and a B, a 5 and a 13_z , and an 11_z ⁴ and an ε . The stock functions as in Pyramid Addition, and the accumulation of points is similar, with the addition of bonus points for removal of bonus pairs mentioned above.

HEXAGON

The game Hexagon is named for the shape of its tableau. It is based on Kings in the Corner solitaire, although it transforms it greatly. The deck is 100_z cards.⁵

The object is to score points by placing cards on the tableau, causing pairs of cards on the same axis (vertical or horizontal) to form a product ending in a factor of a dozen: 2, 3, 4, 6, or 0 (representing 10_z), and optionally 1. So $7 \times 9 = 53_z$ is valid, but $7 \times 3 = 19_z$ is not. All products must not exceed 100_z , precluding $10_z \times 11_z$, $16_z \times 9$, etc.

Face cards, which have no numeric value, automatically go from the face-down center pile to their home piles, marked J, C, B, T, Q, and K. One face card goes into


² *Omega* means one fewer than the base.

³ *Alpha* means one more than the base.

⁴ The author pronounces 11_z as “anzeen,” hence taking the article “an.”

⁵ A decimal version, Pentagon, exists for 100_d physical cards but is not available as an electronic game.

play at a time, causing any face-down cards needed for the play to turn face up. The player then has three ways to create or improve valid scores on the axes surrounding that face card.



In the tableau above, the T is in play. Both products surrounding it are valid: $7 \times 10_z = 70_z$ and $9 \times 6 = 46_z$, totaling 116_z . Switching the position of the 6 and 10 results in $7 \times 6 = 36_z$ and $9 \times 10_z = 90_z$, totaling 126_z , a better score by 10_z . Best is to perform the switch and then place a 15_z card on the 7, for $86_z + 90_z = 176_z$.

If no moves result in two valid products surrounding the face card in play, an existing discard may be used, or a card turned over from the stock into the discard pile.

Points accumulate rapidly in Hexagon, but several kinds of error, such as trying to place a card that makes an invalid product, cause them to be deducted as a penalty. A special effect awaits the player who scores 6,000_z points in a game.

CONCLUSION

Dozenal card games are a viable area of both research and practice, as well as an entertaining tool for teaching and learning arithmetic. Even if not every game is well suited to use in dozenal or with larger decks of cards, the many that are validate both kinds of transformation.

Note: Although I designed the games described, the coding and visual aspects were produced by Thomas Cassidy and Rodrigo Flores. Thanks go also to Jean-Louis Cazaux for making the electronic versions of his K6T cards available. ❏

TIME (AND TIME AGAIN)

by Paul Rapoport

IN THE PREVIOUS ISSUE of the *Duodecimal Bulletin* (WN 33_z),¹ I described some possibilities for dozenal timekeeping, including on a wristwatch face. The face that an expert coder produced was for a Pebble watch. The day we finished the software for it was the day Pebble went out of business, in December 1200_z (2016_d).

At that point I decided to find different watch hardware. Having used various online clocks and that watch for several years, I didn't want to give up on a wristwatch with dozenal time.

Sure, wristwatches are hardly *de rigueur* these days, because many people just use their mobile phone to get the time. True dozenal time is hard to find on a mobile, however. That means not just taking a traditional electronic timepiece and doing something simple, like changing the numerals for ten, eleven, and twelve to dozenal, and the four hash marks between numerals to five. To find a timepiece that divides the day by powers of a dozen—starting with the day or half-day—you probably need either one of my online clocks or Uncial Clock Deluxe,² created by the *Bulletin*'s editor.

THE BANGLE.JS WATCH

While my Pebble watch continued to work, the software on it became uneditable. In early 1205_z, I found the watch Bangle.js (for javascript), which comes from England. Although figuring it out completely was beyond me, I saw immediately that someone could write javascript for it to do what I wanted. Apple watches were not a possibility, because of their requirements and restrictions.

Thomas Cassidy, who did much other work of this sort for me, produced most of the code for that first version of the Bangle watch. When he left the project, David Schonborn finished it. Later in 1205_z, the Bangle.js version 2 hardware came out, sporting several important improvements. I ported my code to the new watch. The face looked the same, showing the date according to my Holocene calendar (see previous issue), followed by dozenal digital time.³

That should have ended the watch project, but it didn't, because I've always wanted a wristwatch in dozenal time that looks analog. Indeed, the very first dozenal timepiece I had someone create, at the University of Illinois, was a large analog clock whose face I clumsily redrew and whose mechanism was



6855-04-07
917.2

¹See https://dozenal.org/drupal/sites_bck/default/files/DuodecimalBulletinIssue531.pdf.

²See <https://sourceforge.net/projects/uncialclock/files/latest/download>.


³Code for this can be found at <https://banglejs.com/apps/?id=doztime>.

transformed to run at half speed (with two hands only). I don't know whether that was the first truly dozenal physical clock, but I do know it was created in 1183_z (1971_d).

On the Bangle.js 2, I was able to create the software myself for an imitation analog watch face, because I found a model for traditional time by Andreas Rozek that I could alter and combine with my previous work to produce what I wanted. Prof. Rozek not only gave permission to use his javascript, but for my watch face solved a few problems that I couldn't.

The result was the first face on left below. The four hands, from slowest to fastest, are red, yellow, green, white. The white hand, which revolves once every 42_z (50_d) seconds of traditional time, is optional. Touching the left side of the face toggles to show or hide it. I hide it, which also reduces battery energy output.

Although the lack of a colored square next to the 2 and 7 is deliberate, I may end up drawing one for each.



TIME IN BALANCED NOTATION

The three faces to the right all illustrate balanced notation for dozenal time, *balanced* because positive and negative digits are given equal prominence. For arithmetic and measurement, balanced notation was discussed and promoted by J. Halcro Johnston in his 1155_z (1937_d) book *The Reverse Notation*.

With balanced time notation, each digit is understood as modifying the previous digit regardless of signs. Obviously, the time 542_z means 500_z trices after midnight plus 40_z trices after that plus 2 trices after that (equivalent to 10:41:40_d). Conversely, the time 5̄42_z means 500_z trices after midnight minus 40_z trices before that plus 2 trices after that (equivalent to 09:21:40_d). The time 5̄4̄2_z means 500_z trices before midnight plus 40_z trices after that minus 2 trices before that (equivalent to 14:38:20_d).

Any time with an initial negative number implies a preceding 1, e.g. 5̄42_z = 15̄42_z. But the 1 is usually omitted for convenience.

With a little practice, all the times are understood without any conversion. Whether balanced notation is better than regular is nonetheless a matter for further discussion.

This idea already has some use in traditional clock time. We often say “twenty to” an hour instead of “forty minutes past” the previous hour, e.g. “20_d to 5” instead of “4:40_d.” The *to* expression clearly indicates a subtraction: 5 o'clock minus 20_d minutes.

Expanding and regularizing that possibility lead to a different but consistent way to read all the hands. When a hand passes the midpoint—the 6—it bumps up the reading of the next slower hand by 1 and itself is indicated as negative. In the example on the far right, regular dozenal time, with digits 0 to 9, would be 516, because each

of the hands is at or past its respective numeral. In balanced notation, that time is 526. The green hand, now on the negative half of the face, increases the yellow hand's number from 1 to 2. Similarly, on the face to its left, the regular dozenal time would be 461; in balanced notation it looks like 561, although when half a trice goes by, the time becomes 562, because then it's closer to that than to 561.

It may be observed that the number read for a hand position in balanced notation is the numeral the hand is closest to, not always the numeral the hand has passed. The two readings are the same until the hand passes the halfway point between two numerals on the face. In that sense, the time with three hands is rounded digitally to the nearest trice, or with four hands to the nearest lull.

In regular dozenal, the third face from the right would read 438, in balanced notation 444.


The three balanced-notation faces are superficial variants of each other. The only difference from regular dozenal notation is that the numerals from 7 to Ɛ are replaced by -5 to -1, with the colored numerals simply representing negatives. I prefer the face with those, because it's less cluttered.

The numerals 6 and 0 are each positive or negative. The 6 is positive for a hand approaching it and negative for a hand having passed it. The 0 has the opposite attributes.

We may also reconfigure the watch face to indicate the day's midnight beginning, uniformly at the bottom here, not with 0 but with -6, and its end, in the same place, with (+)6. That would make noon, at the top, 0. The day would still have 10_z dwells, running from -6 to +6 instead of from 0 to 10_z.

The imitation analog faces make the Holocene calendar date available for a short time by touching their right half, with toggling between cardinal and ordinal date format.

The code for all four faces for the Bangle.js 2 is the same. The user may alter the javascript very slightly to install whichever is wanted. Eventually I may make the choice easier than that. Meanwhile, the code is available to be altered for non-commercial purposes in any way the user prefers. ❏



Motto of the DSA since its founding

by *Paul Rapoport*

THIS CALCULATOR, produced for the Dozenal Society of America, works in both the dozenal and decimal bases and converts between them. Using mostly the Math.js library, it is web-based and thus independent of computer operating system or browser, working on desktop and mobiles. Its usual maximum precision is 20_z (24_d) fractional digits; for complex numbers it is 12_z (14_d), in each part.

Users may set spacing of numbers to 3 digits (e.g. 123 456.789 780 123) or 4 digits (e.g. 1234 5678.9780 1234). There is a full range of trigonometric functions, using radians, degrees, turns (from Primel¹), or unciaPis (from TGM²). Exponents and logarithms to any base are included. Various number theoretical functions are available, including permutations and combinations with and without repetition, factor count ($\tau(n)$), and Euler totient function ($\phi(n)$).

Another unique feature is conversion of measurements among five metrologies: Primel, TGM, USC, Imperial, and Metric. For various measures, e.g. time, velocity, distance, mass, power, electrical current, the calculator converts a measurement from any one metrology to any or all of the others. Measurements from Primel and TGM appear in dozenal, from the others in decimal.

The calculator may be operated at <https://doz-calc.mx-dev.com/>. A button there also brings up the guide to its use.

Comments may be made to paul@rapoport.ca or contact@dozenal.org. Be advised that this calculator may be revised based on user feedback. Users who would like to help us recoup the cost of development and continued maintenance can make a donation to the Dozenal Society of America. Your contributions would be greatly appreciated.

EXAMPLES

Example 1. This example demonstrates a well-known phenomenon: In any base $b \geq 3$, where the three highest digits are $\chi = b - 3$, $\psi = b - 2$, $\omega = b - 1$, when a numeral $n = 123 \dots \chi \omega_b$ is formed by concatenating all the digits of b in ascending order but omitting ψ , the product $\omega \times n$ will be a numeral consisting of ω repetitions of digit 1. When multiplied further by any single digit, the result is ω repetitions of that digit. In dozenal, $\chi = 9$, $\psi = 7$, $\omega = 8$, so e.g. $8 \times 123456789\mathbb{E}_z \times 7 = 777777777777_z$.

$\mathbb{E}^*123456789\mathbb{E}^*7$

7777777777

| | | | | | |
|-----|-------|---------------|-----------|-------------|----------|
| ⌵ | ☐ | ⊖ | 🇺🇸 | 🇬🇧 | 🌐 |
| doz | spc 0 | dig 5 | sto 1 | sci | 1st |
| mod | rad | n | +/- | swap | clear |
| gcd | frs | $\sigma_0(n)$ | $\tau(n)$ | x^2 | x^y |
| nCr | nPr | avg | ln | \log_{10} | \log_y |
| . | % | (|) | ÷ | i |
| 8 | 9 | z | ε | x | del |
| 4 | 5 | 6 | 7 | - | < > |
| 0 | 1 | 2 | 3 | + | = |

Guide (pdf)

¹See page 16 or <https://tinyurl.com/ypyew9ke>

²<https://tinyurl.com/mv4aznym>

Example 2. This gives the square root of the imaginary number i when multiplied by a negative real number. Although 14_z fractional digits are specified, Math.js produces accuracy only to 12_z for complex numbers, as noted above.

| | | | | | |
|---|-----------|---------------------|-----------|---------------|----------|
| sqrt(-100i) | | | | | |
| 8.597 696 503 Ǝ2 97-8.597 696 503 Ǝ22 97i | | | | | |
| \approx | \square | \ominus | | | |
| doz | spc 3 | dig 14 _z | sto 1 | sci | 1st |
| mod | rad | n | +/- | swap | clear |
| gcd | frs | $\sigma_0(n)$ | $\tau(n)$ | \sqrt{x} | x^y |
| nCr | nPr | avg | ln | \log_{10_z} | \log_y |
| . | % | (|) | \div | i |
| 8 | 9 | Ƴ | Ǝ | x | del |
| 4 | 5 | 6 | 7 | - | < > |
| 0 | 1 | 2 | 3 | + | = |

[Guide \(pdf\)](#)

Examples 3 and 4. These demonstrate the identity $\sinh(x) = (e^x - e^{-x})/2$. The equation assumes values in radians. Here for the hyperbolic sine I've specified the angle in Primel turns ($0.6_z \ominus$) instead. It could also be entered as π radians, 1 Pi (TGM), or 130_z degrees.

| | | | | | |
|-----------|-----------|-----------|-------|--------|-------|
| sinh(.6) | | | | | |
| Ǝ.670 27Ǝ | | | | | |
| \approx | \square | \ominus | | | |
| doz | spc 3 | dig 6 | sto 1 | sci | 2nd |
| mod | \odot | n | +/- | swap | clear |
| sin | cos | tan | csc | sec | cot |
| sinh | cosh | tanh | csch | sech | coth |
| . | % | (|) | \div | i |
| 8 | 9 | Ƴ | Ǝ | x | del |
| 4 | 5 | 6 | 7 | - | < > |
| 0 | 1 | 2 | 3 | + | = |

[Guide \(pdf\)](#)

Example 5. This shows the factors of 1000_z. Although they don't all fit into the answer field, scrolling that field reveals the rest of them.

| | | | | | |
|--|-----------|---------------|--------------|---------------|----------|
| frs(1 000) | | | | | |
| 1,2,3,4,6,8,9,10,14,16,20,23,28,30,40,46,54,60,80,9i | | | | | |
| \approx | \square | \ominus | | | |
| doz | spc 3 | dig 5 | sto 1 | sci | 1st |
| mod | rad | n | +/- | swap | clear |
| gcd | pRfrs | $\sigma_0(n)$ | $\varphi(n)$ | x^2 | x^y |
| nCr | nPr | avg | ln | \log_{10_z} | \log_y |
| . | % | (|) | \div | i |
| 8 | 9 | Ƴ | Ǝ | x | del |
| 4 | 5 | 6 | 7 | - | < > |
| 0 | 1 | 2 | 3 | + | = |

[Guide \(pdf\)](#)

| | | | | | |
|--------------------------|-----------|---------------|-----------|---------------|----------|
| $(e^{\pi} - e^{-\pi})/2$ | | | | | |
| Ǝ.670 27Ǝ | | | | | |
| \approx | \square | \ominus | | | |
| doz | spc 3 | dig 6 | sto 1 | sci | 1st |
| mod | rad | n | +/- | swap | clear |
| gcd | frs | $\sigma_0(n)$ | $\tau(n)$ | x^2 | x^y |
| nCr | nPr | avg | ln | \log_{10_z} | \log_y |
| . | % | (|) | \div | i |
| 8 | 9 | Ƴ | Ǝ | x | del |
| 4 | 5 | 6 | 7 | - | < > |
| 0 | 1 | 2 | 3 | + | = |

[Guide \(pdf\)](#)

Examples 6 and 7. These show conversions for distance (length, height, etc.) in Primel, TGM, USC, Imperial, and SI. Again, Primel and TGM are in dozenal, the other three in decimal:

| | | | | | |
|-------|-----------|--------------------|-------|-------------------------------|------------------|
| b lgℓ | 20 | | | | |
| gf | 7£.74£ 19 | | | | |
| ft | 93 | | | | |
| yd | 31 | | | | |
| m | 28.346 4 | | | | |
| dist | □ | ⊖ | 🇺🇸 | 🇬🇧 | 🌐 |
| doz | spc 3 | dig 5 | sto 1 | ⊖ | 1st |
| mod | rad | n | +/- | swap | clear |
| gcd | frs | σ ₀ (n) | τ(n) | x ² | x ^y |
| nCr | nPr | avg | ln | log _{10_z} | log _y |
| . | % | (|) | ÷ | |
| 8 | 9 | 7 | £ | x | del |
| 4 | 5 | 6 | 7 | - | < > |
| 0 | 1 | 2 | 3 | + | = |

Guide (pdf)

| | | | | | |
|------|-----------|--------------------|-------|-------------------------------|------------------|
| lgℓ | 2 000 | | | | |
| gf | 7£.74£ 19 | | | | |
| in | 1 116 | | | | |
| ft | 93 | | | | |
| m | 28.346 4 | | | | |
| dist | □ | ⊖ | 🇺🇸 | 🇬🇧 | 🌐 |
| doz | spc 3 | dig 5 | sto 1 | ⊖ | 1st |
| mod | rad | n | +/- | swap | clear |
| gcd | frs | σ ₀ (n) | τ(n) | x ² | x ^y |
| nCr | nPr | avg | ln | log _{10_z} | log _y |
| . | % | (|) | ÷ | |
| 8 | 9 | 7 | £ | x | del |
| 4 | 5 | 6 | 7 | - | < > |
| 0 | 1 | 2 | 3 | + | = |

Guide (pdf)

The quantity entered was 20_z □biqua-lengthels; the others then appeared automatically. Then I changed three things: □biqua-lengthels to □lengthels (Primel), feet to inches (USC), and yards to feet (Imperial). Each change in units automatically recalculated the quantity:

20_z □biqua-lengthels → 2000_z □lengthels
 93_d feet → 1,116_d inches
 31_d yards → 93_d feet

Although 5 fractional digits are specified, the value in meters needs only 4 for the exact equivalent. The exact foot and yard equivalents require no fractional digits. The Grafut number is rounded.

Example 8. This example converts times of day:

| | | | | | |
|--------|----------|--------------------|-------|-------------------------------|------------------|
| day | .6069 | | | | |
| hour | 10.116 | | | | |
| hour | 12:05:37 | | | | |
| | | | | | |
| | | | | | |
| d-time | □ | ⊖ | 🇺🇸 | 🇬🇧 | 🌐 |
| doz | spc 0 | dig 5 | sto 1 | ⊖ | 1st |
| mod | rad | n | +/- | swap | clear |
| gcd | frs | σ ₀ (n) | τ(n) | x ² | x ^y |
| nCr | nPr | avg | ln | log _{10_z} | log _y |
| . | % | (|) | ÷ | |
| 8 | 9 | 7 | £ | x | del |
| 4 | 5 | 6 | 7 | - | < > |
| 0 | 1 | 2 | 3 | + | = |

Guide (pdf)

The top number, in Primel, is a dozenal fraction of a day; the next, in TGM, is in dozenal hours (quadquaTims). The equivalent in traditional time, in decimal, is the bottom number. It is actually 12:05:37.5_d, although here traditional time's precision is to the most recent whole second, as it is on many time displays. ❄

PRIMEL METROLOGY, PART II

by John Volan

IN THE LAST ISSUE,¹ I presented an introduction to the Primel metrology (brand mark \square), a coherent, dozenal-metric, day/gravity/water-based system of measurement which I have been developing for a number of years. I covered the basic units of mechanics and thermodynamics. I also introduced the concepts of “quantitel” unit names, metrology brand prefixes, and “colloquial” unit names.

This article begins covering more advanced topics, including reciprocal units and angular mechanics. Future articles will cover yet more advanced topics.

“QUANTITELIC” UNITS INSPIRED BY ISO-31

ISO-31² is a standard published by the International Organization for Standardization (ISO).³ Among many other things, it coined several new English words for reciprocals of certain quantities. This took some terminology that had previously been more ad-hoc, and regularized it:

| ISO-31 COINAGE | EXISTING TERMINOLOGY | TECHNICAL MEANING |
|-------------------------|-----------------------------------|--|
| massic quantity | specific quantity | <i>quantity</i> divided by associated mass |
| volumic quantity | [volumic] quantity density | <i>quantity</i> divided by associated volume |
| areic quantity | surface quantity density | <i>quantity</i> divided by associated area |
| lineic quantity | linear quantity density | <i>quantity</i> divided by associated length |

This scheme takes a quantity term (such as *mass*) and applies a common *-ic* suffix to turn it into its reciprocal (*massic*), which then can act as a modifier on some other *quantity*. For instance, the previous article replaced the term *specific thermal capacity* with *massic heatability*.

Primel extends this notion to the unit system, turning any unit name into its reciprocal by appending the **-ic** suffix. The **-ic** suffix may be abbreviated with a backslash (\backslash) to indicate that the reciprocal “quantitelic” unit acts as a denominator placed “under” whatever quantity follows it (if any). For instance:

| QUANTITY | PRIMEL QUANTITEL | QUANTITELIC RECIPROCAL | RECIP ABBREV | METRIC EQUIVALENT |
|----------|--------------------|------------------------|----------------------------|---|
| length | \square lengthel | \square lengthelic | \square lgℓ \backslash | $\approx 1.2192024384_{\text{d}}$ per cm |
| area | \square areanel | \square areanelic | \square arℓ \backslash | $\approx 1.4864545858_{\text{d}}$ per cm ² |
| volume | \square volumel | \square volumelic | \square vmℓ \backslash | $\approx 1.8122890556_{\text{d}}$ per cm ³ |
| mass | \square massel | \square masselic | \square msℓ \backslash | $\approx 1.8123398011_{\text{d}}$ per gram |

¹https://dozenal.org/drupal/sites_bck/default/files/DuodecimalBulletinIssue531.pdf

²https://en.wikipedia.org/wiki/ISO_31, first published in 1992_d (1170_z), superseded by ISO/IEC 80000 (https://en.wikipedia.org/wiki/ISO/IEC_80000) in 2001_d (1179_z).

³https://en.wikipedia.org/wiki/International_Organization_for_Standardization

For example, the previous article identified the \square masselic-heatabilitel ($\square msl\backslash htbl$) as Primel’s coherent unit of massic heatability. We will see more examples of quantitelic units as we go along.

PRIMEL UNITS OF ANGULAR MECHANICS

Angular mechanics (also known as *rotational mechanics*) is the branch of classical mechanics dealing with objects rotating around a fixed axis.

This introduces *plane angle*, or more generally *angular displacement* (symbolized θ), as a distinct type of physical quantity to be measured. For thousands of years, people have been measuring angles using exclusive tools, such as protractors, compasses, and sextants; and have been expressing angle measurements using distinct units not applicable to any other type of quantity. These include **turns** as well as various subdivisions of turns, such as **degrees**, **minutes**, and **seconds** (of arc), although Primel prefers **uncia-turns**, **bicia-turns**, **tricia-turns**, etc., using Systematic Dozenal Nomenclature (SDN) scaling prefixes.⁴

The Radian. As mentioned in the last article, for purposes of physics, the **radian** (abbreviated *rad*) is the most appropriate choice for a coherent unit of angular displacement. This is defined as an angle which subtends a circular arc of length equal to the radius of the enclosing circle. A full turn is τ radians, where $\tau = 2\pi \approx 6.349416967\text{E}635_z$. So 1 radian is approximately $1.7\text{E}0271443577_z$ uncia-turns or 57.2957795130823_d° .

Primel coins **squaradian** (abbreviated *sr*) for the square of the radian. A synonym for this unit, useful in the context of spherical geometry, is the **steradian** (with the same abbreviation). The steradian is the coherent unit of *solid angle* (symbolized Ω). A **spat** is a “full” solid angle covering the entire space surrounding a given vertex. This is σ steradians, where $\sigma = 2\tau = 4\pi \approx 10.696931713\text{E}067_z$.

Angular Dimensionality. Primel holds that angular displacement is a distinct and sensible physical phenomenon, with an irreducible dimensionality of its own, not commensurate with any other physical quantity. Consequently, the radian constitutes another of Primel’s “mundane realities.”

Surprisingly, this is a controversial position, because the International System of Units (SI)⁵, along with many mainstream mathematicians, consider angles to be *dimensionless* quantities. The radian is actually equated with a pure number, i.e. $1 \text{ rad} = 1$.

This stems from SI’s notion that the measure θ of an angle is the ratio of the length s of its subtended arc to the length r of its radius of rotation:

$$\theta = s/r \qquad s = r\theta \qquad r = s/\theta$$

Length over length yields a dimensionless quantity. But this treatment of angles leads to difficulties and inconsistencies across all the quantities of angular mechanics. Radians inexplicably appear and disappear from equations in ad-hoc ways not explicitly driven by the strict algebraic laws of dimensional analysis.

⁴See page 25.

⁵https://en.wikipedia.org/wiki/International_System_of_Units

This has been so troubling that at least a dozen scientists since 1154_z (1936_d)⁶ have written papers advocating for angle to become a first-class dimension, offering various schemes to reconcile the inconsistencies and change the equations of angular mechanics to render them “dimensionally homogeneous.”

The approach I will use here is a variant of an idea proposed by Jacques Romain in 1176_z (1962_d).⁷ Let the “check” diacritic (̃) act as a kind of operator that may be applied to any quantity q to divide it by 1 radian:

$$\check{q} = \frac{q}{1 \text{ rad}}$$

As a trivial case, if we let $q = 1$ (the pure number one), then $\check{1} = 1 \text{ rad}^{-1}$. Applying ISO-31, we can name this the **radianic** as the reciprocal of the radian.⁸ Consequently, we can dub the “check” diacritic the **radianic operator**.

Based on this definition, the radianic operator follows fundamental laws of algebra. Per the associative law, the radianic of a product is the product of either factor multiplied by the radianic of the other; per the distributive law, the radianic of a sum is the sum of the radianics of each of the terms:

$$\check{p \cdot q} = \check{p} \cdot q = p \cdot \check{q} \qquad \check{p + q} = \check{p} + \check{q}$$

So now let an angle θ have first-class physical dimensionality. It retains its intrinsic value no matter what angular units it is measured with. On the other hand, its radianic $\check{\theta}$ is essentially the dimensionless measure quantity when θ is specifically measured in radians: $\theta = \check{\theta} \text{ rad}$. So what SI, and most mathematicians, have been working with all along is not θ , but actually $\check{\theta}$:

$$\check{\theta} = s/r \qquad s = r\check{\theta} \qquad r = s/\check{\theta} \qquad (\text{SI})$$

But Primel endeavors to work only with true dimensioned angles, so the radianic operators must move elsewhere:

$$\theta = s/\check{r} \qquad s = \check{\theta}\theta \qquad \check{r} = s/\theta \qquad (\text{Primel})$$

What is the significance of \check{r} ? It no longer means simply the *radius* of rotation, i.e., the length from the axis of rotation to any point on the circle. Now it means something subtler.

Radiality and the Radiel. I have come to call this new quantity the *radiality* of the rotation (symbolized \check{r}). It is the ratio of a length (the radius) to an angle (1 radian); but in fact, as shown in the equation $\check{r} = s/\theta$ above, it is the ratio of any arc length s around the circle, to the angle θ it subtends.

⁶See https://en.wikipedia.org/wiki/Radian#Dimensional_analysis for discussion and references.

⁷Romain, Jacques E. (July 1962_d). “Angle as a fourth fundamental quantity”. *Journal of Research of the National Bureau of Standards Section B*. **66B** (3): 97. Freely accessible at https://nvlpubs.nist.gov/nistpubs/jres/66B/jresv66Bn3p97_A1b.pdf. He proposed a bracket notation ($\langle q \rangle$) rather than a diacritic.

⁸Some of the solutions to angular dimensionality have involved introducing rad^{-1} as a constant strategically inserted into the equations of angular mechanics. For instance see: Quincey, Paul (1 April 2016_d). “The range of options for handling plane angle and solid angle within a system of units”. *Metrologia*. **53** (2): 840–845. Quincey proposes using η to symbolize $\check{1}$. I prefer to reserve $\eta = \pi/4 = \pi/2$ as a circle constant for the right angle.

Radiality characterizes the curvature of the circular path taken by a rotating object. The smaller the radiality, the more angular displacement occurs per linear displacement, and thus the greater the curvature. The larger the radiality, the less angular displacement occurs per linear displacement, and thus the less the curvature.

Primel's coherent unit of radiality is the \square radiel,⁹ abbreviated $\square rdl$:

$$\square \text{radiel} = \frac{\square \text{lengthel}}{\text{radian}}$$

This is equivalent to $8.20208\bar{3}_d \text{ mm}\cdot\text{rad}^{-1}$. An appropriate colloquial synonym for this is \square morsel-radiality. Scalings of this unit, along with their colloquial synonyms, include:

| | | | |
|--------------------------|---------------------------|-------------------------------|---|
| \square radiel | = $\square rdl$ | = \square morsel-radiality | = $8.20208\bar{3}_d \text{ mm}\cdot\text{rad}^{-1}$ |
| \square unqua-radial | = $\square u\uparrow rdl$ | = \square hand-radiality | = $98.425_d \text{ mm}\cdot\text{rad}^{-1}$ |
| \square biqua-radial | = $\square b\uparrow rdl$ | = \square ell-radiality | = $1.1811_d \text{ m}\cdot\text{rad}^{-1}$ |
| \square triqua-radial | = $\square t\uparrow rdl$ | = \square habital-radiality | = $14.1732_d \text{ m}\cdot\text{rad}^{-1}$ |
| \square quadqua-radial | = $\square q\uparrow rdl$ | = \square stadial-radiality | = $170.0784_d \text{ m}\cdot\text{rad}^{-1}$ |
| \square pentqua-radial | = $\square p\uparrow rdl$ | = \square dromal-radiality | = $2.0409408_d \text{ km}\cdot\text{rad}^{-1}$ |
| \square hexqua-radial | = $\square h\uparrow rdl$ | = \square itineradiality | = $24.4912896_d \text{ km}\cdot\text{rad}^{-1}$ |

The reciprocal of the \square radiel is the \square radielic, abbreviated $\square rdl\setminus$:

$$\square \text{radielic} = \frac{\text{radian}}{\square \text{lengthel}}$$

This is equivalent to approximately $1.219202438404877_d \text{ rad}\cdot\text{cm}^{-1}$.

The \square radiel and the \square radielic turn out to be useful modifiers which can neatly transform units of linear mechanics into units of rotational mechanics.

The squares of these will also prove useful. The square of the \square radiel is the \square squaradiel, abbreviated $\square sdl$:

$$\square \text{squaradiel} = \square \text{radiel}^2 = \frac{\square \text{lengthel}^2}{\text{radian}^2} = \frac{\square \text{areanel}}{\text{steradian}}$$

This is equivalent to $67.2741710069\bar{4}_d \text{ mm}^2\cdot\text{rad}^{-2}$. A useful synonym for this, in the context of spherical geometry and solid angles, is the \square steradiel.

Scalings of this unit, along with their colloquial synonyms, include:

| | | | |
|-------------------------------|----------------------------|-----------------------------------|---|
| \square squaradiel | = $\square sdl$ | = \square morsel-squaradiality | = $67.2741710069\bar{4}_d \text{ mm}^2\cdot\text{rad}^{-2}$ |
| \square biqua-squaradiel | = $\square b\uparrow sdl$ | = \square hand-squaradiality | = $96.87480625_d \text{ cm}^2\cdot\text{rad}^{-2}$ |
| \square quadqua-squaradiel | = $\square q\uparrow sdl$ | = \square ell-squaradiality | = $1.39499721_d \text{ m}^2\cdot\text{rad}^{-2}$ |
| \square hexqua-squaradiel | = $\square h\uparrow sdl$ | = \square habital-squaradiality | $\approx 2.0087969824_d \text{ a}\cdot\text{rad}^{-2}$ |
| \square octqua-squaradiel | = $\square o\uparrow sdl$ | = \square stadial-squaradiality | = $2.892666214656_d \text{ ha}\cdot\text{rad}^{-2}$ |
| \square decqua-squaradiel | = $\square d\uparrow sdl$ | = \square dromal-squaradiality | = $4.16543934910464_d \text{ km}^2\cdot\text{rad}^{-2}$ |
| \square unnilqua-squaradiel | = $\square un\uparrow sdl$ | = \square itineradiality | = $599.823266271068_d \text{ km}^2\cdot\text{rad}^{-2}$ |

The square of the \square radielic is the \square squaradielic, abbreviated $\square sdl\setminus$:

$$\square \text{squaradielic} = \square \text{radielic}^2 = \frac{\text{radian}^2}{\square \text{lengthel}^2} = \frac{\text{steradian}}{\square \text{areanel}}$$

⁹Contraction of radialitel.

This is equivalent to approximately $1.4864545858124_{\text{d}} \text{ rad}^2\text{cm}^{-2}$. In the context of spherical geometry, a useful synonym for this unit is $\square\text{steradielic}$.⁷

Angular Displacement. The rotational analog for length or linear displacement is of course *angular displacement*. From the equation $\theta = s/\check{r}$ above, we see that angular displacement θ (as defined by Primel) is the ratio of arc length s to the radiality \check{r} of the rotation.

Hence one synonym for Primel's coherent unit of angular displacement is the $\square\text{radielic-lengthel}$, abbreviated $\square rd\ell\backslash lgl$:

$$\square\text{radielic-lengthel} = \frac{\text{radian}}{\square\text{lengthel}} \cdot \square\text{lengthel} = \text{radian}$$

Of course this is just a synonym for the **radian**. Another reasonable synonym is $\square\text{ang-lengthel}$,⁸ abbreviated $\square \angle lgl$. (We can generate even more synonyms by substituting synonyms for **lengthel**, including **displacel**, **distancel**, etc.)

Angular Velocity. The rotational analog for linear velocity is *angular velocity* (symbolized ω), which is the time rate of change of angular displacement.

$$\omega = \frac{d\theta}{dt} = \frac{1}{\check{r}} \frac{ds}{dt} = \frac{v_{\perp}}{\check{r}} \quad v_{\perp} = \frac{ds}{dt} \quad (\text{Primel})$$

where v_{\perp} is the tangential (linear) velocity. Note that, because SI uses $\check{\theta}$ rather than true θ , and radius r rather than radiality \check{r} , its version of angular velocity is $\check{\omega}$ rather than true ω :

$$\check{\omega} = \frac{d\check{\theta}}{dt} = \frac{1}{r} \frac{ds}{dt} = \frac{v_{\perp}}{r} \quad v_{\perp} = \frac{ds}{dt} \quad (\text{SI})$$

This gives $\check{\omega}$ a problematic dimensionality indistinguishable from that of frequency (inverse time). Similar analysis applies when considering angular velocity as a vector in three dimensions:

$$\boldsymbol{\omega} = \frac{\check{\mathbf{r}} \times \mathbf{v}}{\|\check{\mathbf{r}}\|^2} \quad \mathbf{v}_{\perp} = \boldsymbol{\omega} \times \check{\mathbf{r}} \quad (\text{Primel})$$

$$\check{\boldsymbol{\omega}} = \frac{\mathbf{r} \times \mathbf{v}}{\|\mathbf{r}\|^2} \quad \mathbf{v}_{\perp} = \check{\boldsymbol{\omega}} \times \mathbf{r} \quad (\text{SI})$$

Primel's coherent unit of angular velocity is the $\square\text{radielic-velocitel}$, abbreviated $\square rd\ell\backslash vcl$:

$$\square\text{radielic-velocitel} = \frac{\text{radian}}{\square\text{lengthel}} \cdot \frac{\square\text{lengthel}}{\square\text{timel}} = \frac{\text{radian}}{\square\text{timel}}$$

⁷Tom Pendlebury, in his TGM metrology (see https://dozenal.org/drupal/sites_bck/default/files/tgm_0.pdf), introduced prefixes **rada**, **radi**, **quara**, **quari**, as TGM's own version of (respectively) **radiel**, **radielic**, **squaradiel**, **squaradielic**. Except that Pendlebury's forms resemble his scaling prefixes, rather than independent unit names.

⁸Contraction of $\square\text{angular-lengthel}$. I will use **ang** as a contraction for **angular** in synonyms for all the quantitels of angular mechanics.

A reasonable synonym for this is \square **ang-velocitel**, abbreviated $\square \triangleleft vcl$. This is equivalent to $34.56_d \text{ rad}\cdot\text{s}^{-1}$. Scalings of this unit include:

| | | | |
|--|--|---|--|
| \square unqua-ang-velocitel | $= \square u \uparrow \triangleleft vcl$ | $= 10_z \text{ rad}\cdot\text{vibe}^{-1}$ | $= 414.72_d \text{ rad}\cdot\text{s}^{-1}$ |
| \square ang-velocitel | $= \square \triangleleft vcl$ | $= 1 \text{ rad}\cdot\text{vibe}^{-1}$ | $= 34.56_d \text{ rad}\cdot\text{s}^{-1}$ |
| \square uncia-ang-velocitel | $= \square u \downarrow \triangleleft vcl$ | $= 1 \text{ rad}\cdot\text{twinkling}^{-1}$ | $= 2.88_d \text{ rad}\cdot\text{s}^{-1}$ |
| \square bicia-ang-velocitel | $= \square b \downarrow \triangleleft vcl$ | $= 1 \text{ rad}\cdot\text{lull}^{-1}$ | $= 14.4_d \text{ rad}\cdot\text{min}^{-1}$ |
| \square tricia-ang-velocitel | $= \square t \downarrow \triangleleft vcl$ | $= 1 \text{ rad}\cdot\text{trice}^{-1}$ | $= 1.2_d \text{ rad}\cdot\text{min}^{-1}$ |
| \square quadcia-ang-velocitel | $= \square q \downarrow \triangleleft vcl$ | $= 1 \text{ rad}\cdot\text{breather}^{-1}$ | $= 6_d \text{ rad}\cdot\text{hr}^{-1}$ |
| \square pentcia-ang-velocitel | $= \square p \downarrow \triangleleft vcl$ | $= 1 \text{ rad}\cdot\text{dwell}^{-1}$ | $= 12_d \text{ rad}\cdot\text{day}^{-1}$ |
| \square hexcia-ang-velocitel | $= \square h \downarrow \triangleleft vcl$ | $= 1 \text{ rad}\cdot\text{day}^{-1}$ | |

Angular Acceleration. The rotational analog of linear acceleration is *angular acceleration* (symbolized α), which is the time rate of change of angular velocity.

$$\alpha = \frac{d\omega}{dt} = \frac{1}{\check{r}} \frac{d^2 s}{dt^2} = \frac{a_{\perp}}{\check{r}} \quad a_{\perp} = \frac{d^2 s}{dt^2} \quad (\text{Primel})$$

where a_{\perp} is the tangential (linear) acceleration. Note that, because SI uses $\check{\omega}$ rather than true ω , and radius r rather than radiality \check{r} , its version of angular acceleration is $\check{\alpha}$ rather than true α :

$$\check{\alpha} = \frac{d\check{\omega}}{dt} = \frac{1}{r} \frac{d^2 s}{dt^2} = \frac{a_{\perp}}{r} \quad a_{\perp} = \frac{d^2 s}{dt^2} \quad (\text{SI})$$

This gives $\check{\alpha}$ a problematic dimensionality indistinguishable from that of frequency squared (inverse time squared).

Similar analysis applies when considering angular acceleration as a vector in three dimensions:

$$\boldsymbol{\alpha} = \frac{\check{\mathbf{r}} \times \mathbf{a}}{\|\check{\mathbf{r}}\|^2} \quad \mathbf{a}_{\perp} = \boldsymbol{\alpha} \times \check{\mathbf{r}} \quad (\text{Primel})$$

$$\check{\boldsymbol{\alpha}} = \frac{\mathbf{r} \times \mathbf{a}}{\|\mathbf{r}\|^2} \quad \mathbf{a}_{\perp} = \check{\boldsymbol{\alpha}} \times \mathbf{r} \quad (\text{SI})$$

Primel's coherent unit of angular acceleration is the \square **radielic-accelerel**, abbreviated $\square rdl\backslash accel$:

$$\square \text{radielic-accelerel} = \frac{\text{radian}}{\square \text{lengthel}} \cdot \frac{\square \text{lengthel}}{\square \text{timel}^2} = \frac{\text{radian}}{\square \text{timel}^2}$$

A reasonable synonym for this is \square **ang-accelerel**, abbreviated $\square \triangleleft accel$. This is equivalent to $1194.3936_d \text{ rad}\cdot\text{s}^{-2}$.

Angular Mass or Moment of Inertia. The rotational analog of mass is known as *moment of inertia* or *angular mass*. SI defines this as the mass m of the rotating object times the square of the radius r of rotation. But Primel uses the radiality \check{r} instead:

$$I = mr^2 \quad (\text{SI}) \quad \check{I} = m\check{r}^2 \quad (\text{Primel})$$

Thus Primel's definition of angular mass is actually SI's version with the radianic operator applied *twice*.

Primel's coherent unit of angular mass is the \square **squaradiel-massel**, abbreviated \square *sdl·msl*:

$$\square\text{squaradiel-massel} = \frac{\square\text{lengthel}^2}{\text{radian}^2} \cdot \square\text{massel}$$

A reasonable synonym for this is \square **ang-massel**, abbreviated \square *∠msl*. This is equivalent to approximately $0.371200648827305_{\text{d}} \text{ g}\cdot\text{cm}^2\text{rad}^{-2}$. A colloquial synonym for this unit is the \square **morsel-ang-mass**.

A useful scaling of this unit is the \square **pentqua-ang-massel**, which may be called colloquially the \square **hand-ang-mass**, because:

$$\begin{aligned} \square\text{hand-ang-mass} &= \square\text{hand-squaradiality} \cdot \square\text{hand-mass} \\ &= \square\text{biqua-squaradiel} \cdot \square\text{triqua-massel} \\ &= \square\text{pentqua-squaradiel-massel} \\ &= \square\text{pentqua-ang-massel} \end{aligned}$$

This is equivalent to approximately $0.923665998489959_{\text{d}} \text{ kg}\cdot\text{dm}^2\text{rad}^{-2}$.

Angular Momentum. The rotational analog of momentum is known as *angular momentum*. SI symbolizes this as **L** and calculates it as the cross-product of the radius vector **r** (position vector of the rotating object relative to the axis of rotation) times the linear momentum vector **p**. SI can also calculate it as the product of its version of angular mass *I* multiplied by its version of angular velocity vector $\check{\omega}$ (sans radians). This is by direct analogy with linear momentum being the product of mass *m* and linear velocity vector **v**:

$$\mathbf{L} = I\check{\omega} = \mathbf{r} \times \mathbf{p} \qquad \mathbf{p} = m\mathbf{v} \qquad (\text{SI})$$

However, this gives **L** a problematic dimensionality indistinguishable from action (which is linear momentum times linear displacement).

In contrast, Primel symbolizes its version of angular momentum as $\check{\check{L}}$ and calculates it as the cross product of the *radiality* vector of the rotating object multiplied by its linear momentum vector **p**. Primel can also calculate it as the product of its squaradianic version of angular mass $\check{\check{I}}$, times its true version of angular velocity vector ω (with radians):

$$\check{\check{L}} = \check{\check{I}}\omega = \check{\check{\mathbf{r}}} \times \mathbf{p} \qquad \mathbf{p} = m\mathbf{v} \qquad (\text{Primel})$$

This gives $\check{\check{L}}$ its own unique dimensionality equivalent to action per angular displacement, analogous to linear momentum being equivalent to action per linear displacement.

This distinction between action and angular momentum is at the heart of why the Planck constant seems to have two values: *h* and $\check{h} = h/\tau$. Both of these are expressed in units of action, but in reality physicists should be using $\check{\check{h}} = \check{h}/\text{rad} = h/(\tau \text{ rad}) = h/\text{turn}$. In other words, when we include the missing angular units, both *h* and \check{h} express the *same* constant: $\check{\check{h}}$, the quantum of angular momentum.

Primel's coherent unit of angular momentum is the \square **radial-momel**,¹⁰ abbreviated \square *rdl·mmℓ*:

$$\begin{aligned}\square\text{radial-momel} &= \frac{\square\text{lengthel} \cdot \square\text{massel} \cdot \square\text{lengthel}}{\text{radian} \quad \square\text{timel}} \\ &= \frac{\square\text{massel} \cdot \square\text{lengthel}^2}{\square\text{timel} \cdot \text{radian}} = \frac{\square\text{actionel}}{\text{radian}}\end{aligned}$$

A reasonable synonym for this is \square **ang-momel**, abbreviated \square *∠mmℓ*.

$$\begin{aligned}\square\text{ang-momel} &= \square\text{ang-massel} \cdot \square\text{ang-velocitel} \\ &= \square\text{squaradiel-massel} \cdot \square\text{radielic-velocitel} \\ &= \square\text{radial-momel}\end{aligned}$$

This is equivalent to approximately $12.8286944234717_{\text{d}} \text{ g}\cdot\text{cm}^2\cdot\text{s}^{-1}\cdot\text{rad}^{-1}$. A colloquial synonym for this unit is \square **morsel-ang-momentum**.

A useful scaling of this unit is the \square **quadqua-ang-momel**, which may be called colloquially the \square **hand-ang-momentum**, because:

$$\begin{aligned}\square\text{hand-ang-momentum} &= \square\text{hand-radiality} \cdot \square\text{hand-momentum} \\ &= \square\text{unqua-radiel} \cdot \square\text{triqua-momel} \\ &= \square\text{quadqua-radiel-momel} \\ &= \square\text{quadqua-ang-momel}\end{aligned}$$

This is equivalent to approximately $2.66015807565108_{\text{d}} \text{ kg}\cdot\text{m}^2\cdot\text{s}^{-1}\cdot\text{rad}^{-1}$.

Angular Force. The rotational analog of force is known as *angular force* or *torque*. SI symbolizes this as \mathbf{T} ,¹¹ and calculates it as the cross-product of the radius vector \mathbf{r} (position vector of the rotating object relative to the axis of rotation) times the linear force vector \mathbf{F} . SI can also calculate it as the product of its version of angular mass I multiplied by its version of angular acceleration vector $\check{\alpha}$ (sans radians). This is by direct analogy with linear force being the product of mass m and linear acceleration vector \mathbf{a} :

$$\mathbf{T} = I\check{\alpha} = \mathbf{r} \times \mathbf{F} \qquad \mathbf{F} = m\mathbf{a} \qquad (\text{SI})$$

However, this gives \mathbf{T} a problematic dimensionality indistinguishable from work (which is linear force times linear displacement).

In contrast, Primel symbolizes its version of angular force as $\check{\check{\mathbf{T}}}$ and calculates it as the cross product of the *radiality* vector of the rotating object multiplied by its linear force vector \mathbf{F} . Primel can also calculate it as the product of its squaradianic version of angular mass $\check{\check{I}}$, times its true version of angular acceleration vector $\check{\alpha}$ (with radians):

$$\check{\check{\mathbf{T}}} = \check{\check{I}}\check{\alpha} = \check{\mathbf{r}} \times \mathbf{F} \qquad \mathbf{F} = m\mathbf{a} \qquad (\text{Primel})$$

¹⁰Recall that Primel's coherent unit of linear momentum, \square **momentumel**, can be contracted to \square **momel**.

¹¹This is uppercase Greek tau, indistinguishable from Latin T. SI actually uses lowercase tau (τ) for this. But I am avoiding this in order to reserve $\tau = 2\pi$ as a circle constant.

This gives $\check{\mathbf{T}}$ its own unique dimensionality equivalent to work (i.e., energy) per angular displacement, analogous to linear force being equivalent to work per linear displacement.

Primel's coherent unit of angular force is the \square **radial-force**, abbreviated $\square rdl.fcl$:

$$\begin{aligned}\square \text{radial-force} &= \frac{\square \text{lengthel}}{\text{radian}} \cdot \frac{\square \text{massel} \cdot \square \text{lengthel}}{\square \text{timel}^2} \\ &= \frac{\square \text{massel} \cdot \square \text{lengthel}^2}{\square \text{timel}^2 \cdot \text{radian}} = \frac{\square \text{workel}}{\text{radian}}\end{aligned}$$

A reasonable synonym for this is \square **ang-force**, abbreviated $\square \angle fcl$.

$$\begin{aligned}\square \text{ang-force} &= \square \text{ang-massel} \cdot \square \text{ang-accelerel} \\ &= \square \text{squaradiel-massel} \cdot \square \text{radielic-accelerel} \\ &= \square \text{radial-force}\end{aligned}$$

This is equivalent to approximately $443.359679275181_d \text{ erg}\cdot\text{rad}^{-1}$ or $44.3359679275181_d \mu\text{J}\cdot\text{rad}^{-1}$.

One useful scaling is the \square **quadqua-radial-force**, which can have the colloquial synonym \square **hand-ang-force**, because:

$$\begin{aligned}\square \text{hand-ang-force} &= \square \text{hand-radiality} \cdot \square \text{hand-force} \\ &= \square \text{unqua-radial} \cdot \square \text{triqua-force} \\ &= \square \text{quadqua-radial-force} \\ &= \square \text{quadqua-ang-force}\end{aligned}$$

This is equivalent to approximately $0.919350630945015_d \text{ J}\cdot\text{rad}^{-1}$.

Translational and Rotational Kinetic Energy. Primel's versions of angular mass (\check{I}) and angular velocity (ω) can be used to calculate the rotational component of an object's kinetic energy (E_R). This is directly analogous to how the translational component of kinetic energy (E_T) can be calculated from the object's mass (m) and linear velocity (v):

$$E_R = \frac{1}{2} \check{I} \omega^2 \qquad E_T = \frac{1}{2} m v^2$$

We can confirm this by looking at Primel's units for these equalities:

$$\begin{aligned}\square \text{energel} &= \square \text{squaradiel-massel} \cdot \square \text{radielic-velocitel}^2 \\ &= (\square \text{radial}^2 \cdot \square \text{massel}) (\square \text{radial}^{-2} \cdot \square \text{velocitel}^2) \\ &= \square \text{massel} \cdot \square \text{velocitel}^2 = \square \text{energel}\end{aligned}$$

Reusability of Unit Names. All of the quantitel and quantitelic unit names described above have been specific to the Primel metrology, so they have all sported Primel's brand mark \square . Nevertheless, they can be reused for other metrologies by simply

| PRIMEL ☐ SUMMARY OF ANGULAR MECHANICAL UNITS | | | | |
|--|---|-----------------------|--|--|
| QUANTITY | QUANTITEL | ABBREV | DECOMPOSITION | METRIC EQUIVALENTS |
| Plane Angle | radian | <i>rad</i> | <i>rad</i> | 1 rad |
| Angular Displacement | ☐radielic-lengthel ☐ang-lengthel | ☐rdl\lgℓ ☐∠lgℓ | | |
| Square Angle | squaradian | <i>sr</i> | <i>rad</i> ² | 1 rad ² |
| Solid Angle | steradian | | | 1 sr |
| Angular Area | ☐squaradielic-areanel ☐steradielic-areanel ☐ang-areanel | ☐sdl\arℓ ☐∠arℓ | | |
| Radiality | ☐radiel | ☐rdℓ | $\frac{\text{☐lgℓ}}{\text{rad}}$ | 8.202083 _d $\frac{\text{mm}}{\text{rad}}$ |
| Inverse Radiality | ☐radielic | ☐rdℓ\ | $\frac{\text{rad}}{\text{☐lgℓ}}$ | ≈ 1.219202438404877 _d $\frac{\text{rad}}{\text{cm}}$ |
| Squaradiality | ☐squaradiel | ☐sdl | $\frac{\text{☐lgℓ}^2}{\text{rad}^2}$ | 67.2741710069 _d $\frac{\text{mm}^2}{\text{rad}^2}$ |
| Steradiality | ☐steradiel | | | |
| Inverse Squaradiality | ☐squaradielic | ☐sdl\ | $\frac{\text{rad}^2}{\text{☐lgℓ}^2}$ | ≈ 1.4864545858124 _d $\frac{\text{rad}^2}{\text{cm}^2}$ |
| Inverse Steradiality | ☐steradielic | | | |
| Angular Velocity | ☐radielic-velocitel ☐ang-velocitel | ☐rdℓ\vcl ☐∠vcl | $\frac{\text{rad}}{\text{☐tml}}$ | 34.56 _d $\frac{\text{rad}}{\text{s}}$ |
| Angular Acceleration | ☐radielic-accelerel ☐ang-accelerel | ☐rdℓ\accl ☐∠accl | $\frac{\text{rad}}{\text{☐tml}^2}$ | 1194.3936 _d $\frac{\text{rad}}{\text{s}^2}$ |
| Angular Mass | ☐squaradiel-massel | ☐sdl·msℓ | $\frac{\text{☐msℓ} \cdot \text{☐lgℓ}^2}{\text{rad}^2}$ | ≈ 0.371200648827305 _d $\frac{\text{g} \cdot \text{cm}^2}{\text{rad}^2}$ |
| Moment of Inertia | | ☐msℓ | | |
| Angular Momentum | ☐radiel-momel ☐ang-momel | ☐rdℓ·mml ☐∠mml | $\frac{\text{☐msℓ} \cdot \text{☐lgℓ}^2}{\text{☐tml} \cdot \text{rad}}$ | ≈ 12.8286944234717 _d $\frac{\text{g} \cdot \text{cm}^2}{\text{s} \cdot \text{rad}}$ |
| Angular Force | ☐radiel-forcel | ☐rdℓ·fcl | $\frac{\text{☐msℓ} \cdot \text{☐lgℓ}^2}{\text{☐tml}^2 \cdot \text{rad}}$ | ≈ 44.3359679275181 _d $\frac{\mu\text{J}}{\text{rad}}$ |
| Torque | ☐ang-forcel | ☐∠fcl | | |

replacing Primel’s brand mark with the brand mark of another metrology (and of course recomputing the unit sizes in terms of that metrology’s own “mundane realities”).

ANGULAR TRANSCENDENTAL FUNCTIONS

An additional problem encountered when considering the dimensionality of angular displacement is the question of how to treat trigonometric and other functions that take angular displacements as arguments. Such functions often have Taylor series expansions, which are infinite sums of terms each containing differing powers of the argument. Such terms would not be commensurate with each other, and so could not be added together, unless the original argument were dimensionless. To see the solution to this issue, let us first consider how to apply the radianic operator to a mathematical function.

For any unary function f , let us define its radianic \check{f} as equivalent to the original function applied to the radianic of its argument:

$$\check{f}(\theta) = f(\check{\theta}) = f\left(\frac{\theta}{1 \text{ rad}}\right)$$

In this way, we have introduced a new “complete” function \check{f} which can accept a dimensioned angular quantity θ . But the first step of \check{f} is to extract the dimensionless radianic measure quantity $\check{\theta}$ and pass that on to f . Then f itself acts as an auxiliary function which only accepts a dimensionless argument, but which is thereby free to calculate any terms it likes from it. If the function is transcendental then it is free to do a Taylor series expansion:

$$\begin{aligned} \widetilde{\sin} \theta &= \sin \check{\theta} = \sum_{n=0}^{\infty} (-1)^n \frac{\check{\theta}^{2n+1}}{(2n+1)!} = \check{\theta} - \frac{\check{\theta}^3}{3!} + \frac{\check{\theta}^5}{5!} - \frac{\check{\theta}^7}{7!} + \dots \\ \widetilde{\cos} \theta &= \cos \check{\theta} = \sum_{n=0}^{\infty} (-1)^n \frac{\check{\theta}^{2n}}{(2n)!} = 1 - \frac{\check{\theta}^2}{2!} + \frac{\check{\theta}^4}{4!} - \frac{\check{\theta}^6}{6!} + \dots \\ \widetilde{\exp i} \theta &= \exp i \check{\theta} = e^{i\check{\theta}} = \sum_{n=0}^{\infty} i^n \frac{\check{\theta}^n}{n!} = 1 + i\check{\theta} - \frac{\check{\theta}^2}{2!} - i\frac{\check{\theta}^3}{3!} + \frac{\check{\theta}^4}{4!} + i\frac{\check{\theta}^5}{5!} - \dots \\ &= \left(1 - \frac{\check{\theta}^2}{2!} + \frac{\check{\theta}^4}{4!} - \dots\right) + i\left(\check{\theta} - \frac{\check{\theta}^3}{3!} + \frac{\check{\theta}^5}{5!} - \dots\right) = \cos \check{\theta} + i \sin \check{\theta} = \widetilde{\cos} \theta + i \widetilde{\sin} \theta \end{aligned}$$

The heart of the matter is that transcendental functions like \sin , \cos and $\exp i$ actually compute relationships between pure mathematical abstractions with no regard to any sort of physical manifestation in the real world. Not even the geometric interpretation as angles in the plane is really pertinent to the abstraction (although it can provide intuition to aid in understanding it).

On the other hand, functions such as $\widetilde{\sin}$, $\widetilde{\cos}$, and $\widetilde{\exp i}$ do relate to real physical quantities that can actually be measured. But to further analyze these quantities, we need to strip them of their physical dimensionality and bridge into the world of pure mathematics. This is fine, but it does mean that we have identified two quite different, yet related, kinds of function, applicable to very different contexts. Problems only arise when we try to conflate the two.

MORE TO COME

Designing Primel’s angular mechanics units has required diverging from SI’s approach to the subject, but as it turns out, not in ways that were entirely unprecedented. The next issue will cover Primel’s units for electricity and magnetism. I will examine not only the usual types of quantities encountered by first-year physics and engineering students, but in fact all the various kinds of quantities embodied in famous equations by James Clerk Maxwell and others.¹² Primel’s approach does not in any way change the dimensionality of these quantities. However, in order to derive a balanced system of quantitel unit names for them, it does make interesting changes to the *terminology* applied to electromagnetic quantities. This may actually prove to be the most controversial aspect of the metrology. I hope you will all get a *charge* out of it. ❏

¹²See https://en.wikipedia.org/wiki/Maxwell's_equations.

| [z] • SYSTEMATIC DOZENAL NOMENCLATURE SUMMARY • [z] | | | | | | |
|---|--------|------|------------|----------------------|------------------|------------------|
| N | ROOT | ABBR | MULTIPLIER | RECIPROCAL | POWER PREFIXES | |
| | | | PREFIX | PREFIX | POSITIVE | NEGATIVE |
| | | | $N \times$ | $\frac{1}{N} \times$ | $10^{+N} \times$ | $10^{-N} \times$ |
| 0 | nil | n | nil- | nilinfra- | nilqua- | nilcia- |
| 1 | un | u | uni- | uninfra- | unqua- | uncia- |
| 2 | bi | b | bina- | bininfra- | biqua- | bicia- |
| 3 | tri | t | trina- | trininfra- | triqua- | tricia- |
| 4 | quad | q | quadra- | quadinfra- | quadqua- | quadcia- |
| 5 | pent | p | penta- | pentinfra- | pentqua- | pentcia- |
| 6 | hex | h | hexa- | hexinfra- | hexqua- | hexcia- |
| 7 | sept | s | septa- | septinfra- | septqua- | septcia- |
| 8 | oct | o | octa- | octinfra- | octqua- | octcia- |
| 9 | enn | e | ennea- | enninfra- | ennqua- | enncia- |
| ĉ | dec | d | deca- | decinfra- | decqua- | decchia- |
| € | lev | L | leva- | levinfra- | levqua- | levcia- |
| 10 | unnil | un | unnili- | unnilinfra- | unnilqua- | unnilcia- |
| 11 | unun | uu | ununi- | ununinfra- | ununqua- | ununcia- |
| 12 | unbi | ub | unbina- | unbininfra- | unbiqua- | unbicia- |
| 13 | untri | ut | untrina- | untrininfra- | untriqua- | untricia- |
| 14 | unquad | uq | unquadra- | unquadinfra- | unquadqua- | unquadcia- |
| 15 | unpent | up | unpenta- | unpentinfra- | unpentqua- | unpentcia- |
| 16 | unhex | uh | unhexa- | unhexinfra- | unhexqua- | unhexcia- |
| 17 | unsept | us | unsepta- | unseptinfra- | unseptqua- | unseptcia- |
| 18 | unoct | uo | unocta- | unoctinfra- | unoctqua- | unoctcia- |
| 19 | unenn | ue | unennea- | unenninfra- | unennqua- | unenncia- |
| 1ĉ | undec | ud | undeca- | undecininfra- | undecqua- | undecchia- |
| 1€ | unlev | uL | unleva- | unlevinfra- | unlevqua- | unlevcia- |
| 20 | binil | bn | binili- | binilinfra- | binilqua- | binilcia- |
| | | | | <i>etc...</i> | | |

uncia was Latin for *one twelfth* • retains same meaning • *inch* and *ounce* are English derivatives
Concatenating roots = positional place-value • Suggested pronunciation: -cia = /ʃə/ (“sha”)
Concatenating prefixes = multiplication • mix & match freely • Commutative Law applies
Prefer Unicode abbreviations where supported • ASCII abbreviations for email, text, etc.

| SDN FORM | EXAMPLE | EXAMPLE | ABBREVIATION | |
|----------------------------|-------------------------------|--------------------------|--------------|--------|
| | VALUE [z] | SDN | UNICODE | ASCII |
| Root Form | 46 | quadhex | qh | qh |
| Multiplier Prefix | 46× | quadhexa- | qh• | qh* |
| With Fractional Part | 4.6× | quad.dot.hexa- | q.h• | q.h* |
| Ordinal | 46 th | quadhexal | qhʹ | qhʹ |
| Reciprocal Prefix | $\frac{1}{46} \times$ | quadhexinfra- | qh\ | qh\ |
| Positive Power Prefix | $10^{+46} \times$ | quadhexqua- | qh† | qh‡ |
| Negative Power Prefix | $10^{-46} \times$ | quadhexcia- | qh‡ | qh# |
| Rational Number | $4 \times \frac{1}{5} \times$ | quadra.pentinfra- | q•p\ | q*p\ |
| Rational Number | $\frac{1}{5} \times 4 \times$ | pentinfra-quadra- | p\q• | p\q* |
| Scientific Notation | $4 \times 10^{+6} \times$ | quadra.hexqua- | q•h† | q*h‡ |
| With Fractional Part | $4.5 \times 10^{+6} \times$ | quad.dot.penta.hexqua- | q.p•h† | q.p*h‡ |
| Scientific Notation | $10^{+6} \times 4 \times$ | hexqua-quadra- | h†q• | h†q* |
| With Fractional Part | $10^{+6} \times 4.5 \times$ | hexqua-quad.dot.penta- | h†q.p• | h†q.p* |
| one dozen years | $10^{+1} \times$ year | unqua-year, unquennium | u†yr | u@yr |
| one gross years | $10^{+2} \times$ year | biqua-year, biquennium | b†yr | b@yr |
| one galore years | $10^{+3} \times$ year | triqua-year, triquennium | t†yr | t@yr |
| two hours (a “dwell”) | $10^{-1} \times$ day | uncia-day | u‡dy | u#dy |
| ten minutes (a “breather”) | $10^{-2} \times$ day | bicia-day | b‡dy | b#dy |
| fifty seconds (a “trice”) | $10^{-3} \times$ day | tricia-day | t‡dy | t#dy |

For more info see:

- Original article: http://www.dozenal.org/drupal/sites_bck/default/files/DSA_kodegadulo_sdn.pdf
- Wiki page: <https://primelmetrology.atlassian.net/wiki/display/PM/Systematic+Numeric+Nomenclature%3A+Dozenal>
- Forum: <https://www.tapatalk.com/groups/dozenonline/systematic-dozenal-nomenclature-f31/>
- Original thread: <https://www.tapatalk.com/groups/dozenonline/systematic-dozenal-nomenclature-t463.html>

AN UNPUBLISHED MANUSCRIPT OF LEIBNIZ'S ON DUODECIMAL



by Lloyd Strickland

MANCHESTER METROPOLITAN UNIVERSITY

lstrickland@mmu.ac.uk



IN THE FIELD OF MATHEMATICS, the German polymath Gottfried Wilhelm Leibniz (1646–1716)¹ is best known for his independent inventions of the calculus and binary arithmetic. Less well known is that Leibniz also invented the base-16 number system, which he called “sedecimal” (see Strickland and Jones 2022). During his exploration of bases 2 and 16, he had occasion to work a little on other number bases, including the duodecimal. While most of his references to base 12 occur in writings that deal with other topics as well, there does exist one manuscript exclusively devoted to it. An English translation of that manuscript is given at the end of this article, the first time the manuscript has been published in any language. By way of an introduction, and to provide some context, I shall begin with an outline of Leibniz’s other engagements with duodecimal, many of which are also unpublished. In contrast to his writings on the binary system, which span hundreds of manuscript pages, Leibniz’s treatment of duodecimal is occasional, scattered, and unsystematic. Yet from those it is clear that Leibniz had a sound understanding of duodecimal, and as we shall see, duodecimal may even have had a role in his invention of binary.

The earliest mention of duodecimal in Leibniz’s writings occurs in a preface he wrote in 1670 to an edition of the writings of the Italian humanist Marius Nizolius (1498–1576). There he considers the view of those who would have it that truth depends upon the definitions of terms, and definitions of terms in turn upon the human mind (in other words, that truth is arbitrary). To this Leibniz (1969, 128) responds: “In arithmetic, and in other disciplines as well, truths remain the same even if notations are changed, and it does not matter whether a decimal or a duodecimal number system is used.” Unfortunately, Leibniz does not reveal where he had learned about duodecimal, though his use of it suggests he thought it was sufficiently well known that educated readers would be able to understand his argument.

Between 1672 and 1676 Leibniz lived in Paris, studying under the tutelage of some of the foremost mathematicians of the day, most notably Christiaan Huygens (1629–1695). As a result of his intense studies he devised the calculus in 1675, and binary arithmetic a few years later. The first references to duodecimal in his mathematical writings date from around this time. For example, in a manuscript entitled “Thesaurus mathematicus” [Mathematical Thesaurus], probably written in either 1678 or 1679, Leibniz works through various topics in arithmetic, geometry, and mechanics; near the end, he outlines how positional notation works in the decimal and then the duodecimal number system:

From this outline it is clear that only these ten digits are needed: 0, 1, 2, 3, 4, 5, 6, 7, 8, 9. Those in the first position signify the equivalent number of 1s, namely no 1s, one 1, two 1s, three, four, five, six, seven, eight, nine 1s. Those in the second position signify the equivalent number of 10s, that is, 1s taken ten times; in the third position, the equivalent number of 100s, that is, 10s taken ten times,

¹ *EDITOR’S NOTE:* All numerals in this article default to decimal [d], unless the text explicitly describes use of duodecimal by Leibniz or his contemporaries. Symbols shown for transdecimal digits are those used by the historical figures themselves.

or the squares of 10; in the fourth position, the equivalent number of *1000s*, that is, *100s* taken ten times, or the cubes of 10, and so on. And in place of 10 one would be able to put any other number, for example, 12. For just as when the base a is 10, the square a^2 signifies 100 and the cube a^3 signifies 1000, so when a is 12, a^2 will be 12 times 12, that is, 144, and a^3 will be 12 times 144. But on this method, instead of the digits mentioned above—0, 1 etc. 9—two new digits would be needed in addition, one which would represent ten, the other which would represent eleven; but [the digits] 10 would signify twelve, and 100 would signify one hundred and forty four. And there are some who prefer to use this method of calculating over the common method, because 12 can be divided by 2, 3, 4, and 6; in addition, a calculation is completed with fewer digits. But the difference is not so great as to be worth abandoning the decimal progression. (LH 35, 1, 25 Bl. 3v)

Leibniz often repeated the claim that some people had preferred duodecimal to decimal, but it is not clear who he had in mind. Late in life he claimed that a proponent of duodecimal had been identified by the German mathematician Daniel Schwenter (1585–1636): “In the German *Deliciae mathematicae*, someone is reported to have given preference to the duodecimal progression, in which eleven digits will be needed, namely 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, δ , ε , where δ is 10 and ε is 11” (LBr 705 Bl. 93r). The reference is probably to Schwenter’s *Deliciae physico-mathematicae* [*The Charms of Physico-Mathematics*] of 1636, which was posthumously revised and expanded by Georg Philipp Harsdörffer (1607–1658) in 1651 and again in 1653. However, as far as I have been able to tell, in none of those works is there any mention of the duodecimal system, let alone any report of anyone endorsing it.² Alternatively, when referring to proponents of duodecimal, Leibniz may have been thinking of Blaise Pascal (1623–1662), who had mentioned duodecimal as an alternative to decimal in an essay presented to the *Académie Parisienne* in 1654 and posthumously published eleven years later. In that essay, Pascal (1665, 42) promised to give a method to determine whether a given number is divisible by any other number, insisting that it would work “not just in our decimal system of numeration (which has been established not as a result of natural necessity, as the common man thinks, but as a result of human custom, and quite foolishly, to be sure), but in a system of numeration based on any progression whatsoever.” To illustrate, he applied his method to the duodecimal system. Leibniz was certainly aware of Pascal’s essay, and may have taken Pascal’s work with duodecimal as an endorsement thereof. He certainly appears to have absorbed what Pascal had to say about duodecimal; in a manuscript written around 1678, Leibniz noted that if the duodecimal system were used, the arithmetic checking method known as casting out nines “could become the proof by casting out elevens” (LH 35, 4, 13 Bl. 21), apparently borrowing the observation from Pascal (1665, 47–48), who had made it some years before.

In the spring of 1680, Leibniz met the Dutch mathematician Johann Jakob Ferguson (1630–1706), and must have mentioned both binary and duodecimal to him, as in the scratch paper upon which both recorded their ideas, Leibniz (1976, 137) wrote out a table showing the values of the decimal numbers 0–8 in binary, and a set of duodecimal digits, with the two extra digits given as \sim and $\$$. In August of the same year, Leibniz noted: “It is well known that all fractions can be expressed by an infinite sequence of integers of a certain progression, for example, the decimal, or even the duodecimal,

²Schwenter (1636, 117–122) does, however, provide a lengthy list of ways that the numbers 2–30 are embedded or reflected in the physical and spiritual realms. With 12, for example, he notes that there were 12 tribes of Israel and 12 apostles, that there are 12 signs of the zodiac, that the year is divided into 12 months etc.

or the one I prefer overall, the binary” (LH 35, 13, 3 Bl. 33). The remainder of this manuscript is concerned with binary fractions, but given the confidence of his remark here, it is likely that he had already undertaken some investigation of duodecimal fractions. If he had, his work on that remains to be discovered.

If Leibniz himself is to be believed, duodecimal even played a role in his invention of binary. The story Leibniz would tell in the 1690s onwards was that he had hit upon binary as the simplest number system from conscious reflection upon the duodecimal and quaternary number systems. In 1697 he wrote:

It is apparent that some considered the duodecimal to be more useful while others took pleasure in the Pythagorean tetractys. At some point it occurred to me to consider what would ultimately be revealed if we used the simplest of all [progressions], namely the dyadic or binary. (Strickland and Lewis 2022, 110)

(The “Pythagorean tetractys,” by the way, is the quaternary—base 4—number system developed by Erhard Weigel (1673).) We should be cautious of at least some of what Leibniz claims here. Certainly, there is no evidence that he knew of Weigel’s quaternary system before 1683, several years after he had invented binary, in which case he could not have been influenced by quaternary. But as we have seen, Leibniz did know about duodecimal at least as far back as 1670, and while the manuscript evidence he left behind does not enable us to verify his later claim that he found his way to binary via duodecimal, it does not enable us to rule it out either.

Although Leibniz wrote relatively little about duodecimal, he was clearly aware of its advantages over decimal, and occasionally indicated that, were the decimal system to be dislodged from common usage, it should be replaced either by the duodecimal or sedecimal. In 1694 or 1695 he wrote:

I think that if anything were to be changed in practice, it would be to use the duodecimal or sedecimal instead of the decimal, for the larger the numbers used by a progression, the more convenient the calculation (Strickland and Lewis 2022, 85).

Leibniz later made the same claim in “Explanation of binary arithmetic,” the only one of his many writings on binary published in his own lifetime. There, after outlining binary notation and arithmetic, Leibniz insisted that binary was not intended to replace decimal in everyday usage because the long strings of digits made it impractical, in which case, he said, it is better to stick with decimal because the numbers are not as long. He then stated: “And if we were accustomed to proceed by twelves or sixteens, there would be even more benefit” (Strickland and Lewis 2022, 196). Despite publicly acknowledging the advantages of duodecimal and sedecimal, Leibniz was no vocal advocate of wholesale reform, nor did he make much use of these number systems in all but a small handful of his extensive mathematical writings.

Let us turn, then, to Leibniz’s unpublished manuscript on duodecimal. It begins by noting that, in the decimal system, the digital root (that is, the digit sum) of multiples of nine is always nine, e.g. $9 \times 3 = 27$, and $2 + 7 = 9$. Leibniz then generalizes this to any number base, or “progression,” supposing that for any base n , the digital root of multiples of $n - 1$ is always $n - 1$. He illustrates this using the duodecimal (or “duodenary”) system, showing that the digital root of any multiple of 11 is always 11, or rather, since he uses the Greek letters χ and ϕ for 10 and 11 respectively, the digital root for any multiple of ϕ is always ϕ . To secure the point, Leibniz draws a table of duodecimal numbers in which his “new notation” for duodecimal is shown alongside the “old meaning” (i.e. the decimal equivalents).

Unfortunately, Leibniz’s motivation for writing the manuscript is unknown—he gives the impression of simply wanting to record an observation he had made, but does not reveal what inspired him to make the observation in the first place. When did he write the piece? The watermark of the manuscript is found in only two of his other writings, one thought to have been written in 1693, the other positively dated to June 1706, which suggests it was written sometime between those two dates. However, it was filed among Leibniz’s mathematical papers of 1695, making it reasonable to think it was written around that time.^{3 4}

REFERENCES

Harsdörffer, Georg Philipp. 1653. *Delitiae mathematicae et physicae der Mathematicchen und Philosophischen Erquickstunden*. Dritter Theil. Nuremberg: Endters.

LBr. = Manuscript held by Gottfried Wilhelm Leibniz Bibliothek – Niedersächsische Landesbibliothek, Hanover, Germany. Cited by shelfmark and Blatt [sheet].

Leibniz, Gottfried Wilhelm. 1969. *Philosophical Papers and Letters*, ed. and trans. Leroy E. Loemker. Dordrecht: D. Reidel.

Leibniz, Gottfried Wilhelm. 1976. *Sämtliche Schriften und Briefe*. Dritte Reihe, Erster Band, ed. Akademie der Wissenschaften der DDR. Berlin: Akademie-Verlag.

LH = Manuscript held by Gottfried Wilhelm Leibniz Bibliothek – Niedersächsische Landesbibliothek, Hanover, Germany. Cited by shelfmark and Blatt [sheet].

Pascal, Blaise. 1665. *Traité du triangle arithmetique, avec quelques autres petits traitez sur la mesme matiere*. Paris: Guillaume Desprez.

Schwenter, Daniel. 1636. *Deliciae physico-mathematicae oder Mathematische und philosophische Erquickstunden*. Nuremberg: Dümmler.

Strickland, Lloyd, and Jones, Owain Daniel. 2022. “F Things You (Probably) Didn’t Know About Hexadecimal”. *The Mathematical Intelligencer*. Available online at: <https://link.springer.com/article/10.1007/s00283-022-10206-w>.

Strickland, Lloyd, and Lewis, Harry. 2022. *Leibniz on Binary: The Invention of Computer Arithmetic*. Cambridge, Mass.: MIT Press.

Weigel, Erhard. 1673. *Tetractys, Summum tum Arithmeticae tum Philosophiae discursivae compendium, artis magnae sciendi*. Jena: Meyer.

³Work on this paper was graciously supported by the Gerda Henkel Stiftung (award AZ 46/V/21).

⁴**EDITOR’S NOTE:** The following pages include a transcription of the original Latin of Leibniz’s manuscript, an English translation on the facing page, and finally an image of the original manuscript itself. The Dozenal Society of America would like to thank Professor Strickland for graciously choosing *The Duodecimal Bulletin* for first publication of this intriguing historical document.

LEIBNIZ: DE PROGRESSIONE DUODENARIA (c. 1695)⁵

Novenarii proprietas est, quae facit ut summa notarum in multiplis ejus novenarium rursus componat:

| | | | | | | | | | | |
|-------|---|----|----|----|----|----|----|----|----|----|
| 9 PER | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| DAT | 9 | 18 | 27 | 36 | 45 | 54 | 63 | 72 | 81 | 90 |

Sed sciendum est hanc proprietatem oriri ex nostro notandi modo qui est arbitrarius; nempe ex eo quod progressionem denariam utimur, et post novem redimus ad 1 adjecta 0. Potuissemus vero alia progressionem quacunque uti et semper haec futura esset proprietatem numeri ultimi in assumpta progressionem. Exempli causa, si pro denaria progressionem adhiberetur duodenaria, ut a quibusdam curiositatis causa factum est, numerus undenarius simili proprietatem gauderet, quod ostendere opera pretium erit. Nempe si duodenaria progressio adhiberetur, numeri decem et undecim proprias acciperent notas, veluti χ pro denario et ϕ pro undenario si placet. Itaque numeri usque ad duodecies duodecim seu centium quadraginta quatuor, ita stabunt:

| | | | | | | | | | | | | |
|-----------------------|----------|----------|----------|----------|----------|----------|----------|----------|----------|------------|------------|----------|
| NOTATIO NOVEM: | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | χ | ϕ | 10 |
| SIGNIFICATIO ANTIQUE: | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 |
| NOTATIO NOVEM: | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 1χ | 1ϕ | 20 |
| SIGNIFICATIO ANTIQUE: | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 |
| NOTATIO NOVEM: | 21 | 22 | 23 | 24 | 25 | 26 | 27 | 28 | 29 | 2χ | 2ϕ | 30 |
| SIGNIFICATIO ANTIQUE: | 25 | 26 | 27 | 28 | 29 | 30 | 31 | 32 | 33 | 34 | 35 | 36 |
| NOTATIO NOVEM: | 31 | 32 | 33 | 34 | 35 | 36 | 37 | 38 | 39 | 3χ | 3ϕ | 40 |
| SIGNIFICATIO ANTIQUE: | 37 | 38 | 39 | 40 | 41 | 42 | 43 | 44 | 45 | 46 | 47 | 48 |
| NOTATIO NOVEM: | 41 | 42 | 43 | 44 | 45 | 46 | 47 | 48 | 49 | 4χ | 4ϕ | 50 |
| SIGNIFICATIO ANTIQUE: | 49 | 50 | 51 | 52 | 53 | 54 | 55 | 56 | 57 | 58 | 59 | 60 |
| NOTATIO NOVEM: | 51 | 52 | 53 | 54 | 55 | 56 | 57 | 58 | 59 | 5χ | 5ϕ | 60 |
| SIGNIFICATIO ANTIQUE: | 61 | 62 | 63 | 64 | 65 | 66 | 67 | 68 | 69 | 70 | 71 | 72 |
| NOTATIO NOVEM: | 61 | 62 | 63 | 64 | 65 | 66 | 67 | 68 | 69 | 6χ | 6ϕ | 70 |
| SIGNIFICATIO ANTIQUE: | 73 | 74 | 75 | 76 | 77 | 78 | 79 | 80 | 81 | 82 | 83 | 84 |
| NOTATIO NOVEM: | 71 | 72 | 73 | 74 | 75 | 76 | 77 | 78 | 79 | 7χ | 7ϕ | 80 |
| SIGNIFICATIO ANTIQUE: | 85 | 86 | 87 | 88 | 89 | 90 | 91 | 92 | 93 | 94 | 95 | 96 |
| NOTATIO NOVEM: | 81 | 82 | 83 | 84 | 85 | 86 | 87 | 88 | 89 | 8χ | 8ϕ | 90 |
| SIGNIFICATIO ANTIQUE: | 97 | 98 | 99 | 100 | 101 | 102 | 103 | 104 | 105 | 106 | 107 | 108 |
| NOTATIO NOVEM: | 91 | 92 | 93 | 94 | 95 | 96 | 97 | 98 | 99 | 9χ | 9ϕ | $\chi 0$ |
| SIGNIFICATIO ANTIQUE: | 109 | 110 | 111 | 112 | 113 | 114 | 115 | 116 | 117 | 118 | 119 | 120 |
| NOTATIO NOVEM: | $\chi 1$ | $\chi 2$ | $\chi 3$ | $\chi 4$ | $\chi 5$ | $\chi 6$ | $\chi 7$ | $\chi 8$ | $\chi 9$ | $\chi\chi$ | $\chi\phi$ | $\phi 0$ |
| SIGNIFICATIO ANTIQUE: | 121 | 122 | 123 | 124 | 125 | 126 | 127 | 128 | 129 | 130 | 131 | 132 |
| NOTATIO NOVEM: | $\phi 1$ | $\phi 2$ | $\phi 3$ | $\phi 4$ | $\phi 5$ | $\phi 6$ | $\phi 7$ | $\phi 8$ | $\phi 9$ | $\phi\chi$ | $\phi\phi$ | 100 |
| SIGNIFICATIO ANTIQUE: | 133 | 134 | 135 | 136 | 137 | 138 | 139 | 140 | 141 | 142 | 143 | 144 |

Hinc jam multipli ipsius undenarii:

| | | | | | | | | | | | | | |
|----------------------------|----|--------|---------|----|----|----|----|----|----|-----|-----|----------|----------|
| id est, communa notatione: | 11 | 22 | 33 | 44 | 55 | 66 | 77 | 88 | 99 | 110 | 121 | 132 | |
| nova notatione erunt: | | ϕ | 1χ | 29 | 38 | 47 | 56 | 65 | 74 | 83 | 92 | $\chi 1$ | $\phi 0$ |

ubi etiam summa binarum notarum semper facit undecim.

⁵ LH 35, 12 1 Bl. 40r. The original Latin.

LEIBNIZ: ON THE DUODENARY PROGRESSION (c. 1695)⁶

It is a property of nines that the sum of the digits in its multiples makes nine again:

| | | | | | | | | | | |
|-------|---|----|----|----|----|----|----|----|----|----|
| 9 BY | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| GIVES | 9 | 18 | 27 | 36 | 45 | 54 | 63 | 72 | 81 | 90 |

But it should be known that this property originates from our way of writing, which is arbitrary, namely, from the fact that we use the decimal progression and after nine we return to 1 by adding 0. But we could use any other progression and the aforementioned property would always be the property of the last digit in the progression adopted. For example, if the duodenary progression were used in place of the decimal, as has been done by some people for the sake of curiosity, the number eleven would enjoy a similar property, which it will be worthwhile to show. Of course, if the duodenary progression were to be used, the numbers ten and eleven would have their own digits, such as χ for ten and ϕ for eleven, if you like. Therefore the numbers up to twelve times twelve, that is, one hundred and forty-four, will be as follows:

| | | | | | | | | | | | | |
|---------------|----------|----------|----------|----------|----------|----------|----------|----------|----------|------------|------------|----------|
| NEW NOTATION: | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | χ | ϕ | 10 |
| OLD MEANING: | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 |
| NEW NOTATION: | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 1χ | 1ϕ | 20 |
| OLD MEANING: | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 |
| NEW NOTATION: | 21 | 22 | 23 | 24 | 25 | 26 | 27 | 28 | 29 | 2χ | 2ϕ | 30 |
| OLD MEANING: | 25 | 26 | 27 | 28 | 29 | 30 | 31 | 32 | 33 | 34 | 35 | 36 |
| NEW NOTATION: | 31 | 32 | 33 | 34 | 35 | 36 | 37 | 38 | 39 | 3χ | 3ϕ | 40 |
| OLD MEANING: | 37 | 38 | 39 | 40 | 41 | 42 | 43 | 44 | 45 | 46 | 47 | 48 |
| NEW NOTATION: | 41 | 42 | 43 | 44 | 45 | 46 | 47 | 48 | 49 | 4χ | 4ϕ | 50 |
| OLD MEANING: | 49 | 50 | 51 | 52 | 53 | 54 | 55 | 56 | 57 | 58 | 59 | 60 |
| NEW NOTATION: | 51 | 52 | 53 | 54 | 55 | 56 | 57 | 58 | 59 | 5χ | 5ϕ | 60 |
| OLD MEANING: | 61 | 62 | 63 | 64 | 65 | 66 | 67 | 68 | 69 | 70 | 71 | 72 |
| NEW NOTATION: | 61 | 62 | 63 | 64 | 65 | 66 | 67 | 68 | 69 | 6χ | 6ϕ | 70 |
| OLD MEANING: | 73 | 74 | 75 | 76 | 77 | 78 | 79 | 80 | 81 | 82 | 83 | 84 |
| NEW NOTATION: | 71 | 72 | 73 | 74 | 75 | 76 | 77 | 78 | 79 | 7χ | 7ϕ | 80 |
| OLD MEANING: | 85 | 86 | 87 | 88 | 89 | 90 | 91 | 92 | 93 | 94 | 95 | 96 |
| NEW NOTATION: | 81 | 82 | 83 | 84 | 85 | 86 | 87 | 88 | 89 | 8χ | 8ϕ | 90 |
| OLD MEANING: | 97 | 98 | 99 | 100 | 101 | 102 | 103 | 104 | 105 | 106 | 107 | 108 |
| NEW NOTATION: | 91 | 92 | 93 | 94 | 95 | 96 | 97 | 98 | 99 | 9χ | 9ϕ | $\chi 0$ |
| OLD MEANING: | 109 | 110 | 111 | 112 | 113 | 114 | 115 | 116 | 117 | 118 | 119 | 120 |
| NEW NOTATION: | $\chi 1$ | $\chi 2$ | $\chi 3$ | $\chi 4$ | $\chi 5$ | $\chi 6$ | $\chi 7$ | $\chi 8$ | $\chi 9$ | $\chi\chi$ | $\chi\phi$ | $\phi 0$ |
| OLD MEANING: | 121 | 122 | 123 | 124 | 125 | 126 | 127 | 128 | 129 | 130 | 131 | 132 |
| NEW NOTATION: | $\phi 1$ | $\phi 2$ | $\phi 3$ | $\phi 4$ | $\phi 5$ | $\phi 6$ | $\phi 7$ | $\phi 8$ | $\phi 9$ | $\phi\chi$ | $\phi\phi$ | 100 |
| OLD MEANING: | 133 | 134 | 135 | 136 | 137 | 138 | 139 | 140 | 141 | 142 | 143 | 144 |

Hence now the multiples of eleven:

| | | | | | | | | | | | | |
|----------------------------------|--------|---------|----|----|----|----|----|----|----|-----|----------|----------|
| that is, in the common notation: | 11 | 22 | 33 | 44 | 55 | 66 | 77 | 88 | 99 | 110 | 121 | 132 |
| will be in the new notation: | ϕ | 1χ | 29 | 38 | 47 | 56 | 65 | 74 | 83 | 92 | $\chi 1$ | $\phi 0$ |

where also the sum of the two digits always makes eleven.

⁶ LH 35, 12 1 Bl. 40r. Translated from the Latin.



DOZENAL MAGIC SQUARES

by Jay L. Schiffman

INTRODUCTORY DISCUSSION

A MAGIC SQUARE is a configuration such that the sum of the elements in every row, every column and along both diagonals is the same. This common sum is referred to as the *magic sum*. For example, if one places each of the first nine counting integers in the following array, a magic square is obtained:

| | | |
|---|---|---|
| 4 | 3 | 8 |
| 9 | 5 | 1 |
| 2 | 7 | 6 |

A natural question one might ask is how the integers were placed in their respective cells. Observe that:

$$\sum_{i=1}^9 i = 1 + 2 + 3 + 4 + 5 + 6 + 7 + 8 + 9 = 45_d = 39_z$$

using Gauss' method (or directly), and $45_d = 39_z$ must be apportioned equally among the three rows, which implied that the sum of the entries in each row, column, and diagonal is $\frac{45}{3}_d = \frac{39}{3}_z = 15_d = 13_z$.¹

Our initial goal hence is to determine which number can be placed in the center square which encompasses four sums; for the entry in the (2,2) position must occur in four different sums that total $15_d = 13_z$; namely from the second row, second column and both diagonals. The only candidate is 5. Observe that

$$15_d = 13_z = 1 + 5 + 9 = 2 + 5 + 8 = 3 + 5 + 7 = 4 + 5 + 6$$

We next focus on the entries situated on the diagonal corners. Such entries occur in three different sums that total $15_d = 13_z$. The entry in the (1,1) position must occur in sums from the first row, first column and the main diagonal. Similarly the entry in the (3,3) position occurs in sums from the third row, third column and the main diagonal. Likewise, the entry in the (1,3) position occurs in sums from the first row, third column and the off diagonal while the entry in the (3,1) position occurs in sums from the third row, first column and the off diagonal. The only possibilities for numbers in the diagonal corners are 2, 4, 6 and 8. Observe that

¹For those readers who are unfamiliar with Gauss' Method, he wrote the sum of the first nine consecutive integers in the first row in the natural order. In the second row, he wrote the sum in reverse order. Adding the two rows, he obtained twice the sum he desired. One would notice that down every column, a sum of ten is obtained, and we have nine addends of ten for a sum of ninety (seven dozen six). Hence the desired sum is half of ninety (seven dozen six) or forty-five (three dozen nine).

$$\begin{aligned}
15_d &= 13_z = 2 + 4 + 9 = 2 + 5 + 8 = 2 + 6 + 7 \\
15_d &= 13_z = 4 + 2 + 9 = 4 + 3 + 8 = 4 + 5 + 6 \\
15_d &= 13_z = 6 + 1 + 8 = 6 + 2 + 7 = 6 + 4 + 5 \\
15_d &= 13_z = 8 + 1 + 6 = 8 + 2 + 5 = 8 + 3 + 4
\end{aligned}$$

Thus if 4 is placed in the (1,1) position, then 6 must be placed in the (3,3) position. Meanwhile if 8 is placed in the (1,3) position, then 2 must be placed in the (3,1) position. This leaves the integers 1, 3, 7 and 9 to be placed in the other cells that are not in the center cell or in the cells on the diagonal corners. These integers are addends in only two different sums. The entry in the (1,2) position occurs in sums from the first row and second column. The entry in the (2,1) position occurs in sums from the second row and first column. The entry in the (2,3) position occurs in sums from the second row and third column. Finally the entry in the (3,2) position occurs in sums from the third row and second column. We note that

$$\begin{aligned}
15_d &= 13_z = 1 + 5 + 6 = 1 + 6 + 8 \\
15_d &= 13_z = 3 + 4 + 8 = 3 + 5 + 7 \\
15_d &= 13_z = 7 + 2 + 6 = 7 + 3 + 5 \\
15_d &= 13_z = 9 + 1 + 4 = 9 + 2 + 4
\end{aligned}$$

Hence if we place 3 in the (1,2) position, then we must place 7 in the (3,2) position. Finally if we place 9 in the (2,1) position, we must place 1 in the the (2,3) position. This completes the magic square.

With the above information, we immerse in the following activities:

1. First, we rotate the magic square above 90_d° clockwise and 90_d° counterclockwise. We next rotate the magic square above both 180_d° clockwise and 180_d° counterclockwise. Finally we illustrate the magic squares obtained and ask if they are really different magic squares.
2. In addition, we add $10_d = 7_z$ to each entry in the original magic square and generate some conclusions.
3. Our final activity in this section involves multiplying each entry in the original magic square by 5 and form some conclusions.

SOLUTIONS TO THE ABOVE IMMERSIONS

1. If we rotate the original magic square

| | | |
|---|---|---|
| 4 | 3 | 8 |
| 9 | 5 | 1 |
| 2 | 7 | 6 |

90_d° clockwise, we obtain the following magic square:

| | | |
|---|---|---|
| 2 | 9 | 4 |
| 7 | 5 | 3 |
| 6 | 1 | 8 |

If we rotate the original magic square 90_d° counterclockwise, we obtain the following magic square:

| | | |
|---|---|---|
| 8 | 1 | 6 |
| 3 | 5 | 7 |
| 4 | 9 | 2 |

If we rotate the original magic square 180_d° counterclockwise, we obtain the following magic square:

| | | |
|---|---|---|
| 6 | 7 | 2 |
| 1 | 5 | 9 |
| 8 | 3 | 4 |

These magic squares are not different from the original one as rotations preserve the magic sum.

2. If we add $10_d = 7_z$ to each entry in the original magic square, we still obtain a magic square with magic sum $45_d = 39_z$:

| | | |
|-----------------|----|-----------------|
| [d] all decimal | | [z] all dozenal |
| 14 | 13 | 18 |
| 19 | 15 | 11 |
| 12 | 17 | 16 |
| | ≡ | 12 |
| | | 11 |
| | | E |
| | | 10 |
| | | 15 |
| | | 14 |

3. If we multiply each entry in the original magic square by 5, we likewise obtain a magic square whose magic sum is five times that of the original magic square and hence $75_d = 63_z$:

| | | |
|-----------------|----|-----------------|
| [d] all decimal | | [z] all dozenal |
| 20 | 15 | 40 |
| 45 | 25 | 5 |
| 10 | 35 | 30 |
| | ≡ | 18 |
| | | 13 |
| | | 34 |
| | | 39 |
| | | 31 |
| | | 5 |
| | | C |
| | | 2E |
| | | 26 |

It is easy to see that if one adds the same constant to each entry in a magic square, the resulting configuration will likewise result in a magic square. To justify this in the (3) case, consider the following magic square with generic entries $a, b, c, d, e, f, g, h, i$ in the configuration below:

| | | |
|-----|-----|-----|
| a | b | c |
| d | e | f |
| g | h | i |

Suppose the sum of the entries in every row, column and both diagonals in this magic square is k . In the above configuration, we arrive at the following eight equations:

$$\begin{array}{lll} a + b + c = k & d + e + f = k & g + h + i = k \\ a + d + g = k & b + e + h = k & c + f + i = k \\ a + e + i = k & g + e + c = k & \end{array}$$

Suppose we add m to each of the entries, we obtain the following magic square with magic sum $k + 3m$.

| | | |
|---------|---------|---------|
| $a + m$ | $b + m$ | $c + m$ |
| $d + m$ | $e + m$ | $f + m$ |
| $g + m$ | $h + m$ | $i + m$ |

RUDOLF ORDREJKA'S MAGIC SQUARE

We first show that the following (3×3) configuration is a magic square consisting of nine primes (in red) attributed to Rudolf Ordrejka (1928-2001)_d (1148-1179)_z with magic sum $177_d = 129_z$:

| | | | | | | | | | | | | | | | | | | | | |
|--|----|-----------------|----|-----|----|---|----|----|-----|--|--|----|----|----|----|----|---|----|----|----|
| [d] all decimal | ≡ | [z] all dozenal | | | | | | | | | | | | | | | | | | |
| <table style="border-collapse: collapse; margin: auto;"> <tr><td style="padding: 2px 10px;">17</td><td style="padding: 2px 10px;">89</td><td style="padding: 2px 10px;">71</td></tr> <tr><td style="padding: 2px 10px;">113</td><td style="padding: 2px 10px;">59</td><td style="padding: 2px 10px;">5</td></tr> <tr><td style="padding: 2px 10px;">47</td><td style="padding: 2px 10px;">29</td><td style="padding: 2px 10px;">101</td></tr> </table> | 17 | 89 | 71 | 113 | 59 | 5 | 47 | 29 | 101 | | <table style="border-collapse: collapse; margin: auto;"> <tr><td style="padding: 2px 10px;">15</td><td style="padding: 2px 10px;">75</td><td style="padding: 2px 10px;">5E</td></tr> <tr><td style="padding: 2px 10px;">95</td><td style="padding: 2px 10px;">4E</td><td style="padding: 2px 10px;">5</td></tr> <tr><td style="padding: 2px 10px;">3E</td><td style="padding: 2px 10px;">25</td><td style="padding: 2px 10px;">85</td></tr> </table> | 15 | 75 | 5E | 95 | 4E | 5 | 3E | 25 | 85 |
| 17 | 89 | 71 | | | | | | | | | | | | | | | | | | |
| 113 | 59 | 5 | | | | | | | | | | | | | | | | | | |
| 47 | 29 | 101 | | | | | | | | | | | | | | | | | | |
| 15 | 75 | 5E | | | | | | | | | | | | | | | | | | |
| 95 | 4E | 5 | | | | | | | | | | | | | | | | | | |
| 3E | 25 | 85 | | | | | | | | | | | | | | | | | | |

Observe the following:

| | |
|-----------------------|----------------------|
| [d] all decimal | [z] all dozenal |
| $17 + 89 + 71 = 177$ | $15 + 75 + 5E = 129$ |
| $113 + 59 + 5 = 177$ | $95 + 4E + 5 = 129$ |
| $47 + 29 + 101 = 177$ | $3E + 25 + 85 = 129$ |
| $17 + 113 + 47 = 177$ | $15 + 95 + 3E = 129$ |
| $89 + 59 + 29 = 177$ | $75 + 4E + 25 = 129$ |
| $71 + 5 + 101 = 177$ | $5E + 5 + 85 = 129$ |
| $17 + 59 + 101 = 177$ | $15 + 4E + 85 = 129$ |
| $47 + 59 + 71 = 177$ | $3E + 4E + 5E = 129$ |

Decimal Case: Note that the units' digits for primes larger than 5 in decimal must necessarily end in one of the digits 1, 3, 7 or 9. Hence adding the constant $\dots 0_d$ (i.e., any integer with any number of decimal digits ending in 0) to every term in the magic square would have the entry in the (2,3) position ending in the digit 5 and hence not prime. Adding the constant $\dots 2_d$ to every term in the magic square would force the entry in the (2,1) position to terminate in the digit 5. Adding the constant $\dots 4_d$ to each term in the magic square would force the entries in the (1,3) and (3,3) positions to end in the digit 5; while adding the constant $\dots 6_d$ to each of

the entries in the second column would yield the resulting entries terminating in the digit 5. Similarly, adding the constant $\dots 8_d$ forces the entries in the (1,1) and (3,1) positions to terminate in the digit 5. Since all the entries in the magic square of primes are odd, adding any integer terminating in an odd digit makes the resulting integer even and thus not prime. Hence it is impossible to add any constant to the magic square to produce a new magic square such that all nine entries are primes.

Dozenal Case: Note that the units' digit of odd primes can end in any one of the dozenal digits 1, 5, 7 or 8. On the other hand, observe that adding the constant $\dots 4_z$ (i.e., an integer with any number of dozenal digits ending in 4) to every term in the original magic square would force the entries in the (1,3), (2,2), and (3,1) positions to terminate in the digit 3 while the entries in each of the other positions to terminate in the digit 9. Thus none of the entries in the new magic square would be prime.

On the other hand, there is the potential to have up to eight of the entries being prime when a positive even integer constant is added to each of the entries in the given magic square of primes. We next cite some examples below with the prime entries in **Red**.

It should be noted that adding any positive odd integer to Rudolph Ordrejka's original magic square would force all of the resulting entries even and larger than two and hence not prime. While infinitely many solutions are generated in this manner, the solution is neither interesting nor elegant.

On the other hand, adding 4 to each entry produces a magic square having no primes and magic sum $189_d = 139_z$:

$$\begin{array}{|c|c|c|} \hline [d] \text{ all decimal} & & \\ \hline 21 & 93 & 75 \\ \hline 117 & 63 & 9 \\ \hline 51 & 33 & 105 \\ \hline \end{array} = \begin{array}{|c|c|c|} \hline [z] \text{ all dozenal} & & \\ \hline 19 & 79 & 63 \\ \hline 99 & 53 & 9 \\ \hline 43 & 29 & 89 \\ \hline \end{array}$$

Each of the entries in this new magic square is divisible by 3. Recall that any dozenal numeral ending in the digits 0, 3, 6 or 9 is divisible by 3.

Moreover if one adds any integer of the form $6n - 2$ ($n \in \mathbb{N}$) to every entry in Rudolf Ordrejka's magic square, one produces a magic square consisting of no primes; for every element in the magic square will be divisible by 3, as evidenced by the digit root decimal, or the final digit in dozenal. Hence adding 4, $10_d = 7_z$, $16_d = 14_z$, etc. to each entry of the original magic square will produce a magic square consisting of no prime entries. Adding 4 to each entry of Rudolf Ordrejka's magic square produces this result for the initial time.

One related observation is to note that every integer in the original magic square has a remainder of two upon division by three. Using the language of congruences in mathematical parlance, we assert that each entry is congruent to two modulo three. In general, we say that a is congruent to b modulo n , written $a \equiv b \pmod{n}$ if and only if a and b have the same remainder upon division by n , or equivalently, $n | (a - b)$.

Hence we observe that $17_d = 15_z \equiv 2 \pmod{3}$; $3 | (17_d - 2 = 15_z - 2) \Leftrightarrow 3 | (15_d = 13_z)$ (since of course $15_d = 13_z = 3 \cdot 5$). The other entries are verified similarly.

We next focus on securing the smallest constants one needs to add to each entry in Ordrejka's prime magic square to produce dozenal magic squares consisting of from

0-8 prime entries inclusive. In each case, we are starting with Rudolph Ordrejka's original magic square, produced below for ready reference.

| | | | | | | | | | | | | | | | | | | | | |
|---|----|-----------------|----|-----|----|---|----|----|-----|---|---|----|----|----|----|----|---|----|----|----|
| [d] all decimal | | [z] all dozenal | | | | | | | | | | | | | | | | | | |
| <table style="width: 100%; border-collapse: collapse;"> <tr><td style="padding: 2px 10px;">17</td><td style="padding: 2px 10px;">89</td><td style="padding: 2px 10px;">71</td></tr> <tr><td style="padding: 2px 10px;">113</td><td style="padding: 2px 10px;">59</td><td style="padding: 2px 10px;">5</td></tr> <tr><td style="padding: 2px 10px;">47</td><td style="padding: 2px 10px;">29</td><td style="padding: 2px 10px;">101</td></tr> </table> | 17 | 89 | 71 | 113 | 59 | 5 | 47 | 29 | 101 | ≡ | <table style="width: 100%; border-collapse: collapse;"> <tr><td style="padding: 2px 10px;">15</td><td style="padding: 2px 10px;">75</td><td style="padding: 2px 10px;">5E</td></tr> <tr><td style="padding: 2px 10px;">95</td><td style="padding: 2px 10px;">4E</td><td style="padding: 2px 10px;">5</td></tr> <tr><td style="padding: 2px 10px;">3E</td><td style="padding: 2px 10px;">25</td><td style="padding: 2px 10px;">85</td></tr> </table> | 15 | 75 | 5E | 95 | 4E | 5 | 3E | 25 | 85 |
| 17 | 89 | 71 | | | | | | | | | | | | | | | | | | |
| 113 | 59 | 5 | | | | | | | | | | | | | | | | | | |
| 47 | 29 | 101 | | | | | | | | | | | | | | | | | | |
| 15 | 75 | 5E | | | | | | | | | | | | | | | | | | |
| 95 | 4E | 5 | | | | | | | | | | | | | | | | | | |
| 3E | 25 | 85 | | | | | | | | | | | | | | | | | | |

Adding $666_d=476_z$ to each entry produces a magic square having one prime for the first time with magic sum $2175_d=1313_z$:

| | | | | | | | | | | | | | | | | | | | | |
|---|-----|-----------------|-----|-----|-----|-----|-----|-----|-----|---|---|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| [d] all decimal | | [z] all dozenal | | | | | | | | | | | | | | | | | | |
| <table style="width: 100%; border-collapse: collapse;"> <tr><td style="padding: 2px 10px;">683</td><td style="padding: 2px 10px;">755</td><td style="padding: 2px 10px;">737</td></tr> <tr><td style="padding: 2px 10px;">779</td><td style="padding: 2px 10px;">725</td><td style="padding: 2px 10px;">671</td></tr> <tr><td style="padding: 2px 10px;">713</td><td style="padding: 2px 10px;">695</td><td style="padding: 2px 10px;">767</td></tr> </table> | 683 | 755 | 737 | 779 | 725 | 671 | 713 | 695 | 767 | ≡ | <table style="width: 100%; border-collapse: collapse;"> <tr><td style="padding: 2px 10px;">48E</td><td style="padding: 2px 10px;">52E</td><td style="padding: 2px 10px;">515</td></tr> <tr><td style="padding: 2px 10px;">54E</td><td style="padding: 2px 10px;">505</td><td style="padding: 2px 10px;">4E5</td></tr> <tr><td style="padding: 2px 10px;">47E</td><td style="padding: 2px 10px;">4E5</td><td style="padding: 2px 10px;">49E</td></tr> </table> | 48E | 52E | 515 | 54E | 505 | 4E5 | 47E | 4E5 | 49E |
| 683 | 755 | 737 | | | | | | | | | | | | | | | | | | |
| 779 | 725 | 671 | | | | | | | | | | | | | | | | | | |
| 713 | 695 | 767 | | | | | | | | | | | | | | | | | | |
| 48E | 52E | 515 | | | | | | | | | | | | | | | | | | |
| 54E | 505 | 4E5 | | | | | | | | | | | | | | | | | | |
| 47E | 4E5 | 49E | | | | | | | | | | | | | | | | | | |

Adding $116_d=98_z$ to each entry produces a magic square having two primes for the first time with magic sum $525_d=379_z$:

| | | | | | | | | | | | | | | | | | | | | |
|---|-----|-----------------|-----|-----|-----|-----|-----|-----|-----|---|---|----|-----|-----|-----|-----|----|-----|-----|-----|
| [d] all decimal | | [z] all dozenal | | | | | | | | | | | | | | | | | | |
| <table style="width: 100%; border-collapse: collapse;"> <tr><td style="padding: 2px 10px;">133</td><td style="padding: 2px 10px;">205</td><td style="padding: 2px 10px;">187</td></tr> <tr><td style="padding: 2px 10px;">229</td><td style="padding: 2px 10px;">175</td><td style="padding: 2px 10px;">121</td></tr> <tr><td style="padding: 2px 10px;">163</td><td style="padding: 2px 10px;">145</td><td style="padding: 2px 10px;">217</td></tr> </table> | 133 | 205 | 187 | 229 | 175 | 121 | 163 | 145 | 217 | ≡ | <table style="width: 100%; border-collapse: collapse;"> <tr><td style="padding: 2px 10px;">E1</td><td style="padding: 2px 10px;">151</td><td style="padding: 2px 10px;">137</td></tr> <tr><td style="padding: 2px 10px;">171</td><td style="padding: 2px 10px;">127</td><td style="padding: 2px 10px;">71</td></tr> <tr><td style="padding: 2px 10px;">117</td><td style="padding: 2px 10px;">101</td><td style="padding: 2px 10px;">161</td></tr> </table> | E1 | 151 | 137 | 171 | 127 | 71 | 117 | 101 | 161 |
| 133 | 205 | 187 | | | | | | | | | | | | | | | | | | |
| 229 | 175 | 121 | | | | | | | | | | | | | | | | | | |
| 163 | 145 | 217 | | | | | | | | | | | | | | | | | | |
| E1 | 151 | 137 | | | | | | | | | | | | | | | | | | |
| 171 | 127 | 71 | | | | | | | | | | | | | | | | | | |
| 117 | 101 | 161 | | | | | | | | | | | | | | | | | | |

Adding $74_d=62_z$ to each entry produces a magic square having three primes for the first time with magic sum $399_d=293_z$:

| | | | | | | | | | | | | | | | | | | | | |
|---|-----|-----------------|-----|-----|-----|----|-----|-----|-----|---|--|----|-----|-----|-----|----|----|----|----|-----|
| [d] all decimal | | [z] all dozenal | | | | | | | | | | | | | | | | | | |
| <table style="width: 100%; border-collapse: collapse;"> <tr><td style="padding: 2px 10px;">91</td><td style="padding: 2px 10px;">163</td><td style="padding: 2px 10px;">145</td></tr> <tr><td style="padding: 2px 10px;">187</td><td style="padding: 2px 10px;">133</td><td style="padding: 2px 10px;">79</td></tr> <tr><td style="padding: 2px 10px;">121</td><td style="padding: 2px 10px;">103</td><td style="padding: 2px 10px;">175</td></tr> </table> | 91 | 163 | 145 | 187 | 133 | 79 | 121 | 103 | 175 | ≡ | <table style="width: 100%; border-collapse: collapse;"> <tr><td style="padding: 2px 10px;">77</td><td style="padding: 2px 10px;">117</td><td style="padding: 2px 10px;">101</td></tr> <tr><td style="padding: 2px 10px;">137</td><td style="padding: 2px 10px;">E1</td><td style="padding: 2px 10px;">67</td></tr> <tr><td style="padding: 2px 10px;">71</td><td style="padding: 2px 10px;">87</td><td style="padding: 2px 10px;">127</td></tr> </table> | 77 | 117 | 101 | 137 | E1 | 67 | 71 | 87 | 127 |
| 91 | 163 | 145 | | | | | | | | | | | | | | | | | | |
| 187 | 133 | 79 | | | | | | | | | | | | | | | | | | |
| 121 | 103 | 175 | | | | | | | | | | | | | | | | | | |
| 77 | 117 | 101 | | | | | | | | | | | | | | | | | | |
| 137 | E1 | 67 | | | | | | | | | | | | | | | | | | |
| 71 | 87 | 127 | | | | | | | | | | | | | | | | | | |

Adding $32_d=28_z$ to each entry produces a magic square having four primes for the first time with magic sum $273_d=179_z$:

| | | | | | | | | | | | | | | | | | | | | |
|--|-----|-----------------|-----|-----|----|----|----|----|-----|---|---|----|----|----|-----|----|----|----|----|----|
| [d] all decimal | | [z] all dozenal | | | | | | | | | | | | | | | | | | |
| <table style="width: 100%; border-collapse: collapse;"> <tr><td style="padding: 2px 10px;">49</td><td style="padding: 2px 10px;">121</td><td style="padding: 2px 10px;">103</td></tr> <tr><td style="padding: 2px 10px;">145</td><td style="padding: 2px 10px;">91</td><td style="padding: 2px 10px;">37</td></tr> <tr><td style="padding: 2px 10px;">79</td><td style="padding: 2px 10px;">61</td><td style="padding: 2px 10px;">133</td></tr> </table> | 49 | 121 | 103 | 145 | 91 | 37 | 79 | 61 | 133 | ≡ | <table style="width: 100%; border-collapse: collapse;"> <tr><td style="padding: 2px 10px;">41</td><td style="padding: 2px 10px;">71</td><td style="padding: 2px 10px;">87</td></tr> <tr><td style="padding: 2px 10px;">101</td><td style="padding: 2px 10px;">77</td><td style="padding: 2px 10px;">31</td></tr> <tr><td style="padding: 2px 10px;">67</td><td style="padding: 2px 10px;">51</td><td style="padding: 2px 10px;">E1</td></tr> </table> | 41 | 71 | 87 | 101 | 77 | 31 | 67 | 51 | E1 |
| 49 | 121 | 103 | | | | | | | | | | | | | | | | | | |
| 145 | 91 | 37 | | | | | | | | | | | | | | | | | | |
| 79 | 61 | 133 | | | | | | | | | | | | | | | | | | |
| 41 | 71 | 87 | | | | | | | | | | | | | | | | | | |
| 101 | 77 | 31 | | | | | | | | | | | | | | | | | | |
| 67 | 51 | E1 | | | | | | | | | | | | | | | | | | |

Adding $18_d=16_z$ to each entry produces a magic square having five primes for the first time with magic sum $231_d=173_z$:

| | | | | | | | | | | | | | | | | | | | | |
|---|-----|-----------------|----|-----|----|----|----|----|-----|---|--|----|----|----|----|----|----|----|----|----|
| [d] all decimal | | [z] all dozenal | | | | | | | | | | | | | | | | | | |
| <table style="width: 100%; border-collapse: collapse;"> <tr><td style="padding: 2px 10px;">35</td><td style="padding: 2px 10px;">107</td><td style="padding: 2px 10px;">89</td></tr> <tr><td style="padding: 2px 10px;">131</td><td style="padding: 2px 10px;">77</td><td style="padding: 2px 10px;">23</td></tr> <tr><td style="padding: 2px 10px;">65</td><td style="padding: 2px 10px;">47</td><td style="padding: 2px 10px;">119</td></tr> </table> | 35 | 107 | 89 | 131 | 77 | 23 | 65 | 47 | 119 | ≡ | <table style="width: 100%; border-collapse: collapse;"> <tr><td style="padding: 2px 10px;">2E</td><td style="padding: 2px 10px;">8E</td><td style="padding: 2px 10px;">75</td></tr> <tr><td style="padding: 2px 10px;">7E</td><td style="padding: 2px 10px;">65</td><td style="padding: 2px 10px;">1E</td></tr> <tr><td style="padding: 2px 10px;">55</td><td style="padding: 2px 10px;">3E</td><td style="padding: 2px 10px;">9E</td></tr> </table> | 2E | 8E | 75 | 7E | 65 | 1E | 55 | 3E | 9E |
| 35 | 107 | 89 | | | | | | | | | | | | | | | | | | |
| 131 | 77 | 23 | | | | | | | | | | | | | | | | | | |
| 65 | 47 | 119 | | | | | | | | | | | | | | | | | | |
| 2E | 8E | 75 | | | | | | | | | | | | | | | | | | |
| 7E | 65 | 1E | | | | | | | | | | | | | | | | | | |
| 55 | 3E | 9E | | | | | | | | | | | | | | | | | | |

Adding 2 to each entry produces a magic square having six primes for the first time with magic sum $183_d=133_z$:

| | | |
|-----------------|----|-----|
| [d] all decimal | | |
| 19 | 91 | 73 |
| 115 | 61 | 7 |
| 49 | 31 | 103 |

 \equiv

| | | |
|-----------------|----|----|
| [z] all dozenal | | |
| 17 | 77 | 61 |
| 97 | 51 | 7 |
| 41 | 27 | 87 |

Adding $14_d=12_z$ to each entry produces a magic square having seven primes for the first time with magic sum $219_d=163_z$:

| | | |
|-----------------|-----|-----|
| [d] all decimal | | |
| 31 | 103 | 85 |
| 127 | 73 | 19 |
| 61 | 43 | 115 |

 \equiv

| | | |
|-----------------|----|----|
| [z] all dozenal | | |
| 27 | 87 | 71 |
| 77 | 61 | 17 |
| 51 | 37 | 97 |

Adding $12_d=10_z$ to each entry produces a magic square having eight primes for the first time with magic sum $213_d=159_z$:

| | | |
|-----------------|-----|-----|
| [d] all decimal | | |
| 29 | 101 | 83 |
| 125 | 71 | 17 |
| 59 | 41 | 113 |

 \equiv

| | | |
|-----------------|----|----|
| [z] all dozenal | | |
| 25 | 85 | 6E |
| 75 | 5E | 15 |
| 4E | 35 | 95 |

Note that such answers are not unique and the reader can secure multiple solutions! For example, adding $26_d=22_z$ to each entry in Rudolf Ordrejka’s magic square likewise produces a magic square having 6 primes with magic sum $255_d=193_z$, namely:

| | | |
|-----------------|-----|-----|
| [d] all decimal | | |
| 43 | 115 | 97 |
| 139 | 85 | 31 |
| 73 | 55 | 127 |

 \equiv

| | | |
|-----------------|----|----|
| [z] all dozenal | | |
| 37 | 97 | 81 |
| E7 | 71 | 27 |
| 61 | 47 | 77 |

ADDITIONAL OBSERVATIONS

It was earlier mentioned that the magic squares produced by adding any integer of the form $6n - 2$ ($n \in \mathbb{N}$) to each entry of Rudolf Ordrejka’s magic square, one produces a magic square consisting of no primes; for every element in the magic square will be divisible by three by the sum of the digits test. The reason for this follows from the following congruence (for we are adding to elements congruent to two modulo three elements that are congruent to one modulo three so that the sum is congruent to zero modulo three):

$$\text{If } a \equiv b \pmod{n} \text{ and } c \equiv d \pmod{n}, \text{ then } a + c \equiv b + d \pmod{n}. \quad (1)$$

For example:

| [d] all decimal | [z] all dozenal |
|--|--|
| $17 \equiv 2 \pmod{3} \Leftrightarrow$ | $15 \equiv 2 \pmod{3} \Leftrightarrow$ |
| $17 + 4 \equiv (2 + 4) \pmod{3} \Leftrightarrow$ | $15 + 4 \equiv (2 + 4) \pmod{3} \Leftrightarrow$ |
| $21 \equiv 6 \pmod{3} \Leftrightarrow$ | $19 \equiv 6 \pmod{3} \Leftrightarrow$ |
| $3 (21 - 6) \Leftrightarrow$ | $3 (19 - 6) \Leftrightarrow$ |
| $3 15 \Leftrightarrow$ | $3 13 \Leftrightarrow$ |
| $15 = 3 \cdot 5$ | $13 = 3 \cdot 5$ |

To prove (1), note that:

$$\begin{aligned} \text{Since } a &\equiv b \pmod{n}, & n|(a - b) & \quad (2) \\ \text{Since } c &\equiv d \pmod{n}, & n|(c - d) & \quad (3) \end{aligned}$$

Now using elementary properties of divisibility, since $n|(a - b)$ and $n|(c - d)$,

$$\begin{aligned} n|[a - b] + [c - d] &\Leftrightarrow \\ n|[a - b + c - d] &\Leftrightarrow \\ n|[a + c - b - d] &\Leftrightarrow \\ n|[a + c] - [b + d] &\Leftrightarrow \\ a + c &\equiv (b + d) \pmod{n} \end{aligned}$$

An additional question was raised with regards to the above. Is it possible to find a constant to add to Rudolph Ordrejka's original magic square to obtain one consisting of no primes, but none of the entries is divisible by three? The possibility for this certainly exists; for as one proceeds further out in the set of positive primes (which is infinite and first proven by Euclid over two thousand years ago), the gap between consecutive primes can be made as wide as one pleases by considering the following argument in which we seek to secure n consecutive composite integers:

Consider the sequence

$$\{(n + 1)! + 2, (n + 1)! + 3, (n + 1)! + 4, \dots, (n + 1)! + (n + 1)\}$$

The first term is divisible by 2, the second term is divisible by 3, the third term is divisible by 4 and so forth while the last term is divisible by $n + 1$. Hence one has n consecutive composite integers. This sequence is definitely not minimal.

For example, to secure a sequence consisting of five consecutive integers utilizing this constructive process, we form the sequence:

$$\begin{aligned} (5 + 1)! + 2 &= 6! + 2 = 720_d + 2 = 722_d = 500_z + 2 = 502_z \\ (5 + 1)! + 3 &= 6! + 3 = 720_d + 3 = 723_d = 500_z + 3 = 503_z \\ (5 + 1)! + 4 &= 6! + 4 = 720_d + 4 = 724_d = 500_z + 4 = 504_z \\ (5 + 1)! + 5 &= 6! + 5 = 720_d + 5 = 725_d = 500_z + 5 = 505_z \\ (5 + 1)! + 6 &= 6! + 6 = 720_d + 6 = 726_d = 500_z + 6 = 506_z \end{aligned}$$

The first term is divisible by 2, the second term is divisible by 3, the third term is divisible by 4, the fourth term is divisible by 5 and the fifth term is divisible by 6. Of course the minimal sequence of five consecutive composite integers is far smaller, namely $\{24_d, 25_d, 26_d, 27_d, 28_d\} \equiv \{20_z, 21_z, 22_z, 23_z, 24_z\}$.


If one reverts to the previous question, I noted that the difference between the largest and smallest integers in the original magic square is $113_d - 5 = 108_d = 95_z - 5 = 90_z$.

Hence I looked to secure a prime gap of this size. The integer $2238823_d=8\text{E}\text{E}747_z$ is prime and there is a gap of $108_d=90_z$ for the first time following this prime. The next prime is $2238931_d=8\text{E}\text{E}817_z$. Hence the integers from $2238824_d=8\text{E}\text{E}748_z$ to $2238930_d=8\text{E}\text{E}816_z$ are all composite.

If we add the constant $2238822_d=8\text{E}\text{E}746_z$ to all terms of the original magic square, we obtain a magic square consisting of no primes with none of the entries divisible by three.

Is there a minimum constant that achieves our goal? Indeed the answer to this question is in the affirmative and a MATHEMATICA Program was run to accomplish this. It turns out that the constants must be congruent to either zero modulo three or two modulo three so that the resulting sums with the entries (all congruent to two modulo three in the original magic square) are congruent to either two modulo three or one modulo three respectively and hence are not divisible by three. The constants $1656_d = \text{E}60_z \equiv 0(\text{mod } 3)$ and $2876_d = 17\text{E}8_z \equiv 2(\text{mod } 3)$ are optimal.

CONCLUSION

This article served to furnish some engaging mathematics dealing with magic squares that is accessible and hopefully fun mathematics for students and teachers from a wide spectrum of grade levels ranging from elementary through college and beyond. In addition, the posing of thoughtful questions is key for a productive teaching and learning experience. With regards to *The Standards For Mathematical Practice* articulated in The Common Core, this article attempted to demonstrate the necessity to make sense of problems and persevere in solving them, make use of structure, use appropriate tools strategically, construct viable arguments and critique the reasoning of others and express regularity in repeated reasoning (modular arithmetic). One might argue with regards to the other practices as well as the related NCTM works of Peg Smith and Mary Kay Stein. The references below will provide the reader with additional magic squares to partake in as well as related matters. The excellent websites Mathworld and The On-Line Encyclopedia of Integer Sequences can further excite, invigorate and truly fuel one's passion for mathematics. I invite the reader to further partake of these ideas and always enjoy the mathematical journey! 

REFERENCES

1. *MathWorld - A Wolfram Resource*, Champaign-Urbana, IL (2018_d) (1202_z).
2. *Principles and Standards for School Mathematics*, The National Council of teachers of Mathematics (NCTM), Reston, VA (1990_d) (1197_z).
3. *The Standards For Mathematical Practice*, The Common Core, Achieve (2005_d) (11E1_z).
4. Smith, P. and Stein, M.K., *5 Practices for Orchestrating Productive Mathematical Discussions*, Corwin Press, NCTM (2011_d) (11E7_z).
5. *OEIS, The On-Line Encyclopedia of Integer Sequences*, www.oeis.org, ATT Labs, Murray Hill, NJ (2018_d) (1202_z).



POUNDS, SHILLINGS, PENCE

THE COMMONWEALTH'S DOZENAL CURRENCY

by Andon Epp

CURRENCY is not the first application of dozenal that one may think of. Measurement in dozenal has been well explored, as there are many dozens already present in some unit relationships (20_z hours/day, 10_z months/year, 10_z inches/foot, and more), and several attempts to create a coherent dozenal metric system have been devised, some of which were explored in *Bulletin* issue $\zeta 3_z$.¹

Global systems of money, however, have been mostly decimal for a biquennium or two. Starting with the Russian ruble and US dollar, the division of basic units of currency into 100_d parts gradually became standard throughout the 11st_z and 12nd_z biquennia (18th_d through 20th_d centuries). Today, with these subdivisions ingrained into digital bank accounts and credit card data, the chance appears slim that they will be changed, although some countries have eliminated their low-value physical coins. However, some non-decimal systems of money lingered on, and the most notable and interesting for dozenalists was pounds, shillings, and pence (£sd), which predominated in Britain and Commonwealth nations.

The system consisted of one dozen (12_d) pence to a shilling, and one dozen eight (20_d) shillings to a pound. Combinations of amounts were written and spoken in these three units: for example, as three shillings and fivepence or “three and five” (3s 5d or $3/5$), or two pounds, thirteen shillings and sevenpence (£2/13/7_d). It originated from ancient Roman accounting,² which at first recognized the bronze *as* and silver *denarius*, originally in a ten-to-one relationship; later the *solidus* and *libra* were added. The relationships between these units were inconsistent among various regions until the eighth century (sixth biquennium), when the French emperor Charlemagne declared that the *libra* would be valued at a troy pound of silver, worth 20_d *solidi* or 240_d *denarii* (silver pennyweights). This standardization soon spread through continental Europe; it became known as *livres-sous-deniers* in France, and England (along with the rest of Britain) adopted them as pounds-shillings-pence after the Norman conquest. Note that the same abbreviations, £sd, were used for these, regardless of language.

The UK minted coins of various denominations during the use of £sd, including fractions of pence. Table 1^{3 4 5} shows these coins with their values and mintage dates. The dozen-to-one ratio of the shilling and penny made the lower denominations well suited for dozenal numeration.⁶ The circulating coins from farthing to sixpence come out as simple dozenal radix expressions and fractions of a shilling. The system carried with it the benefits of the dozen's divisibility; a shilling could be split

¹https://dozenal.org/drupal/sites_bck/default/files/DuodecimalBulletinIssue531.pdf

²<https://en.wikipedia.org/wiki/%C2%A3sd>

³https://en.wikipedia.org/wiki/List_of_British_banknotes_and_coins

⁴The “halfpenny” was more commonly contracted to “ha’penny.”

⁵The “crown” was used mostly for commemorative purposes.

⁶Covered in A.C. Aitken's “The Case Against Decimalisation,” which also entertained the idea of a “royal,” a dozen-shilling unit. <http://dozenalsociety.org.uk/pdfs/aitken.pdf>

TABLE 1. VALUES AND MINTAGE ERAS OF VARIOUS £SD COINS

| NAME | VALUE [d] | VALUE [z] | MINTAGE [d] | MINTAGE [z] |
|------------------|--------------|-----------------|----------------|----------------|
| Quarter farthing | $1/16$ d | 0.009 s | 1839–1868 | 1093–10E8 |
| Third farthing | $1/12$ d | 0.01 s | 1827–1913 | 1083–1135 |
| Half farthing | $1/8$ d | 0.016 s | 1828–1856 | 1084–1078 |
| Farthing | $1/4$ d | 0.03 s | c.1200–1960 | c.840–1174 |
| Halfpenny | $1/2$ d | 0.06 s | 1260–1969 | 890–1181 |
| Penny | 1 d | 0.1 s | 757–1970 | 531–1182 |
| Thruppenny bit | 3 d | 0.3 ($1/4$) s | 1547–1970 | 78E–1182 |
| Groat | 4 d | 0.4 ($1/3$) s | 1836–1862 | 1090–10E2 |
| Sixpence | 6 d | 0.6 ($1/2$) s | 1547–1970 | 78E–1182 |
| Shilling | 1 s (12 d) | 1 s | 1502–1970 | 752–1182 |
| Florin | 2 s | 2 s | 1848–1970 | 1070–1182 |
| Half crown | 2 s 6 d | 2.6 s | 1528–1969 | 774–1181 |
| Double florin | 4 s | 4 s | 1887–1890 | 1113–1116 |
| Crown | 5 s | 5 s | 1528–1969 | 774–1181 |
| Half sovereign | 10 s | 7 s | 1817–1937 | 1075–1155 |
| Sovereign | £1 (20 s) | 18 s | 1817–1937 | 1075–1155 |

into 2, 3, 4, 6, and even 8 if halfpence were used. The “pence table” learned in schools simply employed dozenal multiplication, with several repeating patterns. There were also slight mental arithmetic advantages: if the price of a dozen items was 3/– (three shillings), each item clearly cost 3d each.

The pound’s value of twenty shillings made £sd a hybrid system, which made working large sums of money more difficult, and it became the main reason the system was abandoned. The British half-crown ($2/6$), at $1/8$ of a pound, functioned much like the US quarter, and its value was even converted to 25_dc in some countries during decimalization. The double florin ($1/5$ of £1) may have accelerated decimal currency had it caught on with the public, but it was only minted for four years. There was even a short-lived “guinea” series based on multiples/fractions of 19_z (21_d) shillings, but the coins were unpopular, although guineas were still used as value terms. In fact, since prices and amounts above around £5 were uncommon prior to post-decimal inflation in the 1180_z (1970_s_d), the dozenal portion of £sd predominated in practice.

The UK and Commonwealth began debating decimal conversion of money in the 1170_z (1960_s_d), with each country soon making the switch in conjunction with its adoption of metric units. Most, such as South Africa, Australia, and New Zealand, chose to make ten former shillings the new unit, dividing it into 100_d cents. This led to a shilling worth 10_dc and a new 1c coin close in value to the old penny. The British government elected to keep the pound, making a shilling 5 “new pence” (100_d p making £1). Coins from the sixpence upward could still be used in transactions at their decimal values after the conversion (see Table 2), but they have since ceased to be legal tender.

Indeed, £sd was not strictly *dozenal*, as it also had a strong *vigesimal* element, and all the multiples of each denomination were expressed in decimal anyway. Consequently,

TABLE 2. CONVERSION OF £SD COINS TO DECIMAL.

| COIN | ABBR [d] | VALUE (UK) in £p [d] | VALUE (AUS, NZ) in \$c [d] |
|-------------------|-------------|-------------------------|-------------------------------|
| Penny | 1d | Approx. ½p | Approx. 1c |
| Thrupenny bit | 3d | Approx. 1p | Approx. 2c |
| Sixpence | 6d | 2½p | 5c |
| Shilling | 1/- | 5p | 10c |
| Florin | 2/- | 10p | 20c |
| Crown | 5/- | 25p | 50c |
| Ten-shilling note | 10/- | 50p (coin) | \$1 |
| Pound note | £1 | £1 (note) | \$2 |

on the DozensOnline forum,⁷ some members of the DSA and DSGB have pondered what true dozenal money could look like in principle.

In a thread titled “What Unit Would You Use in a New Dozenal Currency,”⁸ user “Dan” proposes retaining the main unit (be it the dollar, pound, euro, or anything else) and divide it into uncias. He notes that this would preserve America’s half-dollars and quarters. However, a bicia-dollar would be worth less than the penny, which would be problematic since the penny is already being considered for discontinuation.

In the same thread, user “Joshbuckler” proposes retaining the penny as is, but building up a new main currency unit out of a gross of pennies. User “Kodegadulo” echoes this in the thread “A Problem and a Solution to Converting Decimal to Dozenal Currency.”⁹ This would avoid the issue of rounding error when converting from decimal currency to dozenal. Given how much software there is these days which is keeping track of financial accounts precisely to the penny, such rounding errors could be especially disruptive. That said, replacing a country’s main currency unit with something else runs the risk of stripping its currency of its international prestige.⁷

See Table 3 to compare and contrast these proposals. (Bicia-dollars are included for comparison, but are shown in gray.)

Proposals for coins also differ. The most popular solution is a 1-3-6-10-30-60_z system. This would be analogous to Europe’s 1-2-5-10-20-50_d system, but with the added benefit of each denomination being a multiple of the previous one (see Table 3). However, other configurations, including 1-3-10_z and 1-4-6-10_z, are also viable.

Regularizing currency around twelves alone would make for clean divisibility throughout the system. For instance, dividing any dozenal power by 3 gives 4 of the next power down, and this works at all powers:

$$\begin{aligned} \$0.10_z \div 3 &= \$0.04_z \\ \$1.00_z \div 3 &= \$0.40_z \\ \$10.00_z \div 3 &= \$4.00_z \\ \$100.00_z \div 3 &= \$40.00_z \end{aligned}$$

⁷ <https://www.tapataalk.com/groups/dozensonline/currency-systems-f20/>

⁸ <https://www.tapataalk.com/groups/dozensonline/viewtopic.php?t=598>

⁹ <https://www.tapataalk.com/groups/dozensonline/viewtopic.php?t=2148>

⁷ Kodegadulo himself may not be taking this proposal completely seriously, because he whimsically suggests calling the new bigua-penny unit a U.S. “zachary” (with abbreviation \mathbb{Z}) in honor of Zachary Taylor. Zachary Taylor happens to be the *dozen*th president of the United States—but not a particularly memorable one, since he died after only a year in office, apparently of indigestion.

Kodegadulo also suggests calling the quadqua-penny or bigua-zachary ($\mathbb{Z}100.00_z$) a “woodrow”, after Woodrow Wilson. Wilson happened to be president on July 4, 1140_z (1920_d), the first biquennial anniversary of the Declaration of Independence. However, given that he spent much of that year incapacitated by a stroke, it was not the most auspicious celebration.

Contrast this with £sd which carried the dozen in only one position. A third of a shilling was simply 4 pence, but a third of a pound was 6 shillings 8 pence.

$$1s \div 3 = 4d$$

$$£1 \div 3 = 6s 8d$$

In a decimal world, the pragmatic choice was to enhance decimal with auxiliary factors of three, because they are missing in decimal yet useful enough to warrant inclusion. This can be seen in cases such as sexagesimal time, US customary measures, the ISO 2848 standard for metric building construction,⁸ and of course £sd currency. But in a dozenal world, with fully dozenal measurement and timekeeping, as well as fully dozenal currency, would there be an analogous choice to enhance dozenal with auxiliary factors of five? Such a world would likely feel no need to augment the already-rich divisors of dozenal; a factor of five might seem almost as alien as seven. So the inconveniences of independent arithmetic methods could disappear.

When the United Kingdom completed its program with Decimal Day on February 13_z, 1183_z (February 15_d, 1971_d)¹⁰ a long-lasting and interesting currency system was discontinued. The historical significance of pounds, shillings, and pence, however, has inspired some dozenalists to go all in on using their favorite number to divvy up the bill. ❧

TABLE 3. POTENTIAL CONVERSIONS OF
DECIMAL MONEY TO DOZENAL

| RETAIN THE DOLLAR AND DIVIDE IT DOZENALLY | | RETAIN THE PENNY PROPOSE BIQUA-PENNY AS MAIN UNIT | |
|---|------------------------------|---|------------------------------|
| COIN/NOTE VALUE [z] | EQUIVALENT DOLLARS [d] | COIN/NOTE VALUE [z] | EQUIVALENT DOLLARS [d] |
| \$0.01 | \$0.0069 $\bar{4}$ | ₤0.01 | \$0.01 |
| \$0.03 | \$0.0208 $\bar{3}$ | ₤0.03 | \$0.03 |
| \$0.06 | \$0.041 $\bar{6}$ | ₤0.06 | \$0.06 |
| \$0.10 | \$0.08 $\bar{3}$ | ₤0.10 | \$0.12 |
| \$0.30 | \$0.25 | ₤0.30 | \$0.36 |
| \$0.60 | \$0.50 | ₤0.60 | \$0.72 |
| \$1.00 | \$1.00 | ₤1.00 | \$1.44 |
| \$3.00 | \$3.00 | ₤3.00 | \$4.32 |
| \$6.00 | \$6.00 | ₤6.00 | \$8.64 |
| \$10.00 | \$12.00 | ₤10.00 | \$17.28 |
| \$30.00 | \$36.00 | ₤30.00 | \$51.84 |
| \$60.00 | \$72.00 | ₤60.00 | \$103.68 |
| \$100.00 | \$144.00 | ₤100.00 | \$207.36 |

⁸Which uses “preferred numbers” that include a factor of 3, to introduce 300_d mm (“metric foot”), 600_d mm, and 1200_d mm units. See https://en.wikipedia.org/wiki/ISO_2848

¹⁰For a retrospective on the process of decimalization in the UK, see the “Funny Money” video series: (1) <https://youtu.be/1jeW5LHzNKg>, (2) https://youtu.be/4tnn_8TX1Ec, (3) <https://youtu.be/pC0f2uP26wk>.

POETRY CORNER

MULTIVERSE MULTI-VERSE

COULD THERE BE PARALLEL WORLDS where English-speaking people (including poets) tally their numbers in **twelves** or **eights** or **twenties**, as the case may be? Here's a well-known rhyme by an anonymous Elizabethan bard (NOT!) from a tantalizing triptych of alternate timelines:

DOZENAL-VERSE

*Twenty-six days hath September,
April, June, and November.
Twenty-seven for the rest,
Save Febru'ry doth fail this test,
Having only twenty-four,
Tho' it once possess'd two more.
Whither they went hath long been
reckon'd:
July stole one, August the second.
But when a leap year doth arrive,
'Twill gain one back, for twenty-five.
Every four years mark that date,
Save once in ev'ry tenzy-eight.
Trimming thus the septdubquennia,
Will keep us, for a few triquennia,
Tracking ev'ry passing season
With perfect rhythm, rhyme & reason.*

OCTAL-VERSE

*Throcty-six days hath September,
April, June, and November.
Throcty-seven for the rest,
Save Febru'ry doth fail this test,
Having only throcty-four,
Tho' it once possess'd two more.
Whither they went hath long been
reckon'd:
July stole one, August the second.
But when a leap year doth arrive,
'Twill gain one back, for throcty-five,
Quadrennially earning such reward,
Save once each second checkerboard.
Trimming thus the septdubquennia,
Will keep us, for a few triquennia,
Tracking ev'ry passing season
With perfect rhythm, rhyme & reason.*


VIGESIMAL-VERSE

*Score & ten days hath September,
April, June, and November.
Score & 'leven for the rest,
Save Febru'ry doth fail this test,
Having only eight & score,
Tho' it once possess'd two more.
Whither they went hath long been
reckon'd:
July stole one, August the second.
But when a leap year's stars align,
'Twill gain one back, for score & nine.
Every four years mark that date,
Save once in each six-score & eight.
Trimming thus the septdubquennia,
Will keep us, for a few triquennia,
Tracking ev'ry passing season
With perfect rhythm, rhyme & reason.*

All of these verses imply the same leap-year calculation based on powers of 2, shown below.^{1 2} This yields an average year precisely equal to the nominal tropical year of three hundred sixty-five days, five hours, forty-eight minutes, and forty-five seconds; or **two gross sixty-five days, two dwells, ten breathers, and ten and a half trices**; or **five checkerboard, fivocety-five days and octy-seven-and-a-half percheckerboard**; or **eighteen-score-and-five days and four perscore, sixteen perscoresquare and seventeen-and-a-half perscorecube**. Despite involving no round centuries or biquennia, this scheme more accurately tracks the seasons than the Gregorian calendar average of 365.2425₄ days. "Unleap" years, applying the rule proleptically, are shown at bottom. (Bases indicated by color. Latin uppercase transdecimals used for consistency.)

| DECIMAL | DOZENAL | OCTAL | VIGESIMAL |
|-------------------------------------|---------------------------------------|---------------------------------------|---|
| 30 × 4 = 120 | 26 × 4 = A0 | 36 × 4 = 170 | 1A × 4 = 60 |
| 31 × 7 = 217 | 27 × 7 = 161 | 37 × 7 = 331 | 1B × 7 = AH |
| 28 × 1 = 28 | 24 × 1 = 24 | 34 × 1 = 34 | 18 × 1 = 18 |
| 1 ÷ 4 = 0.25 | 1 ÷ 4 = 0.3 | 1 ÷ 4 = 0.2 | 1 ÷ 4 = 0.5 |
| -1 ÷ 128 = <u>-0.0078125</u> | -1 ÷ A8 = <u>-0.0116</u> | -1 ÷ 200 = <u>-0.004</u> | -1 ÷ 68 = <u>-0.032A</u> |
| 365.2421875 | 265.2AA6 | 555.174 | I5.4GHA |
| DAYS | DAYS | DAYS | DAYS |
| 2 ⁻² = 06:00:00 | 2 × C ⁻¹ = 04:00:00 | 1 × 8 ⁻¹ = 03:00:00 | 4 × K ⁻¹ = 04:48:00.0 |
| -2 ⁻⁷ = <u>-00:11:15</u> | A × C ⁻² = 01:40:00 | 7 × 8 ⁻² = 02:37:30 | G × K ⁻² = 00:57:36.0 |
| 05:48:45 | A × C ⁻³ = 00:08:20 | 4 × 8 ⁻³ = <u>00:11:15</u> | H × K ⁻³ = 00:03:03.6 |
| | 6 × C ⁻⁴ = <u>00:00:25</u> | 05:48:45 | A × K ⁻⁴ = <u>00:00:05.4</u> |
| | 05:48:45 | | 05:48:45.0 |
| UNLEAP YEARS | UNLEAP YEARS | UNLEAP YEARS | UNLEAP YEARS |
| 128 640 1152 1664 | A8 454 800 B68 | 200 1200 2200 3200 | 68 1C0 2HC 434 |
| 256 768 1280 1792 | 194 540 8A8 1054 | 400 1400 2400 3400 | CG 1I8 340 49C |
| 384 896 1408 1920 | 280 628 994 1140 | 600 1600 2600 3600 | J4 24G 3A8 4Q0 |
| 512 1024 1536 2048 | 368 714 A80 1228 | 1000 2000 3000 4000 | 15C 2B4 3GG 528 |

¹In Systematic Numeric Nomenclature (SNN) generalized to multiple bases, *septdubquennium* means 2⁷ years.
²Historically, February did not directly lose days to July and August—but it's a fun conceit.



THE DOZENAL SOCIETY OF AMERICA

APPLICATION FORM

LAST: _____ FIRST: _____ MID.: _____

ADDRESS: _____

CITY: _____ STATE: _____ ZIP: _____

COUNTRY: _____

PHONE(S): _____

EMAIL(S): _____

DEGREES: _____

OTHER SOCIETY MEMBERSHIPS: _____

DESIRED MEMBERSHIP LEVEL (CHECK ONE):

REGULAR (\$12.00_d) SUBSCRIPTION (\$36.00_d) DONATION: _____

A subscription membership entitles you to a paper copy of *The Duodecimal Bulletin* when it is published, for a period of one year from your donation; regular members receive only a digital copy. Donations of any amount, large or small, help keep the DSA going and are greatly appreciated.

To facilitate communication, do you grant the DSA permission to furnish your name and contact information to other DSA members? Yes No

We'd be delighted to see you at our meetings, and are always interested in your thoughts, ideas, and participation. Please tell us about your particular interests here:

Please mail this form with any donation to:

The Dozenal Society of America
604-36 South Washington Square # 815
Philadelphia, PA 19106-4154

Or sign up on our web page:

<http://www.dozenal.org/drupal/content/member-signup>



The Dozenal Society of America
3100 Central Ave, Suite 412
St Petersburg, FL 33712

Founded 1160_z
(1944_d)